

(19)



Europäisches Patentamt
European Patent Office
Office européen des brevets



(11)

EP 0 867 504 A1

(12)

EUROPEAN PATENT APPLICATION

(43) Date of publication:
30.09.1998 Bulletin 1998/40

(51) Int Cl.⁶: **C12N 9/28, C12N 15/56,
C11D 3/386**

(21) Application number: **98109967.4**

(22) Date of filing: **10.02.1994**

(84) Designated Contracting States:
**AT BE CH DE DK ES FR GB GR IE IT LI LU MC NL
PT SE**

(30) Priority: **11.02.1993 US 16395**

(62) Document number(s) of the earlier application(s) in
accordance with Art. 76 EPC:
94909609.3 / 0 689 589

(71) Applicant: **GENENCOR INTERNATIONAL INC.
Rochester, New York 14618 (US)**

(72) Inventors:
• **Solhelm, Leif
Clinton, Iowa 52732 (US)**

- **Power, Scott D.
San Bruno, CA 94066 (US)**
- **Requadt, Carol A.
Tiburon, CA 94920 (US)**
- **Mitchinson, Colln
Half Moon Bay, CA 94019 (US)**

(74) Representative: **Baldock, Sharon Claire et al
BOULT WADE TENNANT,
27 Farnival Street
London EC4A 1PQ (GB)**

Remarks:

This application was filed on 02 - 06 - 1998 as a
divisional application to the application mentioned
under INID code 62.

(54) **Oxidation-stable alpha-amylase**

(57) A mutant alpha-amylase is provided that is the
expression product of a mutated DNA sequence encod-
ing an alpha-amylase, the mutated DNA being derived
from a precursor alpha-amylase, which is a *Bacillus al-*

pha-amylase, by substitution or deletion of an amino
and at a position equivalent to M + 15 in *B licheniformis*
alpha-amylase. The mutant alpha-amylase are suitable
for use in detergent compositions and in process for
starch liquification.

EP 0 867 504 A1

Description

This application is a divisional application from European Patent Application No 94909609.3 filed 10th February 1994.

The present invention relates to novel alpha-amylase mutants having an amino acid sequence not found in nature, such mutants having an amino acid sequence wherein one or more amino acid residue(s) of a precursor alpha-amylase, specifically an oxidizable amino acid, have been substituted with a different amino acid. The mutant enzymes of the present invention exhibit altered stability/activity profiles including but not limited to altered oxidative stability, altered pH performance profile, altered specific activity and/or altered thermostability. In a particular embodiment the invention provides *Bacillus* alpha-amylases having a substitution or deletion of an amino acid at a position equivalent to M + 15 in *Bacillus licheniformis* alpha-amylase and provides uses of these alpha-amylase.

Alpha-amylases (alpha-1,4-glucan-4-glucanohydrolase, EC3.2.1.1) hydrolyze internal alpha-1,4-glucosidic linkages in starch largely at random, to produce smaller molecular weight malto-dextrins. Alpha-amylases are of considerable commercial value, being used in the initial stages (liquefaction) of starch processing; in alcohol production; as cleaning agents in detergent matrices; and in the textile industry for starch desizing. Alpha-amylases are produced by a wide variety of microorganisms including *Bacillus* and *Aspergillus*, with most commercial amylases being produced from bacterial sources such as *B. licheniformis*, *B. amyloliquefaciens*, *B. subtilis*, or *B. stearothermophilus*. In recent years the preferred enzymes in commercial use have been those from *B. licheniformis* because of their heat stability and performance, at least at neutral and mildly alkaline pH's.

Previously there have been studies using recombinant DNA techniques to explore which residues are important for the catalytic activity of amylases and/or to explore the effect of modifying certain amino acids within the active site of various amylases (Vihinen, M. et al. (1990) J. Biochem. **107**:267-272; Holm, L. et al. (1990) Protein Engineering **3**: 181-191; Takase, K. et al. (1992) Biochimica et Biophysica Acta, **1120**:281-288; Matsui, I. et al. (1992) Feds Letters Vol. 310, No. 3, pp. 216-218); which residues are important for thermal stability (Suzuki, Y. et al. (1989) J. Biol. Chem. **264**:18933-18938); and one group has used such methods to introduce mutations at various histidine residues in a *B. licheniformis* amylase, the rationale for making substitutions at histidine residues was that *B. licheniformis* amylase (known to be thermostable) when compared to other similar *Bacillus* amylases, has an excess of histidines and, therefore, it was suggested that replacing a histidine could affect the thermostability of the enzyme (Declerck, N. et al. (1990) J. Biol. Chem. **265**:15481-15488; FR 2 665 178-A1; Joyet, P. et al. (1992) Bio Technology **10**:1579-1583).

It has been found that alpha-amylase is inactivated by hydrogen peroxide and other oxidants at pH's between 4 and 10.5 as described in the examples herein. Commercially, alpha-amylase enzymes can be used under dramatically different conditions such as both high and low pH conditions, depending on the commercial application. For example, alpha-amylases may be used in the liquefaction of starch, a process preferably performed at a low pH (pH < 5.5). On the other hand, amylases may be used in commercial dish care or laundry detergents, which often contain oxidants such as bleach or peracids, and which are used in much more alkaline conditions.

In order to alter the stability or activity profile of amylase enzymes under varying conditions, it has been found that selective replacement, substitution or deletion of oxidizable amino acids, such as methionine, tryptophan, tyrosine, histidine or cysteine, results in an altered profile of the variant enzyme as compared to its precursor. Because currently commercially available amylases are not acceptable (stable) under various conditions, there is a need for an amylase having an altered stability and/or activity profile. This altered stability (oxidative, thermal or pH performance profile) can be achieved while maintaining adequate enzymatic activity, as compared to the wild-type or precursor enzyme. The characteristic affected by introducing such mutations may be a change in oxidative stability while maintaining thermal stability or *vice versa*. Accordingly, the substitution of different amino acids for an oxidizable amino acid(s) in the alpha-amylase precursor sequence or the deletion of one or more oxidizable amino acid(s) may result in altered enzymatic activity at a pH other than that which is considered optimal for the precursor alpha-amylase. In other words, the mutant enzymes of the present invention may also have altered pH performance profiles, which may be due to the enhanced oxidative stability of the enzyme.

The present invention relates to novel alpha-amylase mutants that are the expression product of a mutated DNA sequence encoding an alpha-amylase, the mutated DNA sequence being derived from a precursor alpha-amylase by the deletion or substitution (replacement) of one or more oxidizable amino acid. In particular the invention relates to a mutant alpha-amylase that is the expression product of mutated DNA sequence encoding an alpha-amylase, the mutated DNA sequence being derived from a precursor alpha-amylase which is a *Bacillus* alpha-amylase by substitution or deletion of an amino acid at a position equivalent to M + 15 in *Bacillus licheniformis* alpha-amylase.

In another embodiment of the present invention the mutants comprise a substitution of one or more tryptophan residues alone or in combination with the substitution of one or more methionine residues in a precursor alpha-amylase. Such mutant alpha-amylases, in general, are obtained by *in vitro* modification of a precursor DNA sequence encoding a naturally occurring or recombinant alpha-amylase to encode the substitution or deletion of one or more amino acid residues in a precursor amino acid sequence.

The substitution or deletion of one or more amino acids in the amino acid sequence is due to the replacement or deletion of one or more methionine and/or tryptophan, residues in such sequence. These oxidizable amino acid residues may be replaced by any of the other 20 naturally occurring amino acids. If the desired effect is to alter the stability of the precursor, the amino acid residue may be substituted with a non-oxidizable amino acid (such as alanine, arginine, asparagine, aspartic acid, glutamic acid, glutamine, glycine, isoleucine, leucine, lysine, phenylalanine, proline, serine, threonine, or valine) or another oxidizable amino acid (such as cysteine, methionine, tryptophan, tyrosine or histidine, listed in order of most easily oxidizable to less readily oxidizable). Likewise, if the desired effect is to alter thermostability, any of the other 20 naturally occurring amino acids may be substituted (i.e., cysteine may be substituted for methionine).

The methionine to be replaced is a methionine at a position equivalent to position + 15 in *B. licheniformis* alpha-amylase. The preferred substitute amino acids at position + 15 are leucine (L), threonine (T), asparagine (N), aspartate (D), serine (S), valine (V) and isoleucine (I), although other substitute amino acids not specified above may be useful. A specifically preferred mutant of the present invention is M15L.

Another embodiment of this invention relates to mutants comprising the substitution of a tryptophan residue equivalent to any of the tryptophan residues found in *B. licheniformis* alpha-amylase (see Fig. 2). Preferably the tryptophan to be replaced is at a position equivalent to +138 in *B. licheniformis* alpha-amylase. A mutation (substitution) at a tryptophan residue may be made alone or in combination with mutations at other oxidizable amino acid residues. Specifically, it may be advantageous to modify by substitution of at least one tryptophan in combination with at least one methionine.

The alpha-amylase mutants of the present invention, in general, exhibit altered oxidative stability in the presence of hydrogen peroxide and other oxidants such as bleach or peracids, or, more specific, milder oxidants such as chloramine-T. Mutant enzymes having enhanced oxidative stability will be useful in extending the shelf life and bleach, perborate, percarbonate or peracid compatibility of amylases used in cleaning products. Similarly, reduced oxidative stability may be useful in industrial processes that require the rapid and efficient quenching of enzymatic activity. The mutant enzymes of the present invention may also demonstrate a broadened pH performance profile whereby mutants such as M15L show stability for low pH starch liquefaction. The mutants of the present invention may also have altered thermal stability whereby the mutant may have enhanced stability at either high or low temperatures. It is understood that any change (increase or decrease) in the mutant's enzymatic characteristic(s), as compared to its precursor, may be beneficial depending on the desired end use of the mutant alpha-amylase.

In addition to starch processing and cleaning applications, variant amylases of the present invention may be used in any application in which known amylases are used, for example, variant amylases can be used in textile processing, food processing, etc. Specifically, it is contemplated that a variant enzyme, inactivated by oxidation, would be useful in a process where it is desirable to completely remove amylase activity at the end of the process, for example, in frozen food processing applications.

The preferred alpha-amylase mutants of the present invention are derived from a *Bacillus* strain such as *B. licheniformis*, *B. amyloliquefaciens*, and *B. stearothermophilus*, and most preferably from *Bacillus licheniformis*.

In another aspect of the present invention there is provided a novel form of the alpha-amylase normally produced by *B. licheniformis*. This novel form, designated as the A4 form, has an additional four alanine residues at the N-terminus of the secreted amylase. (Fig. 4b.) Derivatives or mutants of the A4 form of alpha-amylase are encompassed within the present invention. By derivatives or mutants of the A4 form, it is meant that the present invention comprises the A4 form alpha-amylase containing one or more additional mutations such as, for example, mutation (substitution, replacement or deletion) of one or more oxidizable amino acid(s).

In a composition embodiment of the present invention there are provided detergent compositions, liquid, gel or granular, comprising the alpha-amylase mutants described herein. Additionally, it is contemplated that the compositions of the present invention may include an alpha-amylase mutant having more than one site-specific mutation.

In yet another composition embodiment of the present invention there are provided compositions useful in starch processing and particularly starch liquefaction. The starch liquefaction compositions of the present invention preferably comprise an alpha-amylase mutant having a substitution or deletion at position M15. Additionally, it is contemplated that such compositions may comprise additional components as known to those skilled in the art, including, for example, antioxidants, calcium, ions, etc.

In a process aspect of the present invention there are provided methods for liquefying starch, and particularly granular starch slurries, from either a wet or dry milled process. Generally, in the first step of the starch degradation process, the starch slurry is gelatinized by heating at a relatively high temperature (up to about 110°C). After the starch slurry is gelatinized it is liquefied and dextrinized using an alpha-amylase. The conditions for such liquefaction are described in US patent applications 07/785,624 and 07/785, 623 and US Patent 5,180,699. The present method for liquefying starch comprises adding to a starch slurry an effective amount of an alpha-amylase of the present invention, alone or in combination with additional excipients such as an antioxidant, and reacting the slurry for an appropriate time and temperature to liquefy the starch.

A further aspect of the present invention comprises the DNA encoding the mutant alpha-amylases of the present

invention (including A4 form and mutants thereof) and expression vectors encoding the DNA as well as host cells transformed with such expression vectors

The invention will now be described by way of example with reference to the accompanying drawings:-

Fig. 1 shows the DNA sequence of the gene for alpha-amylase from *B. licheniformis* (NCIB8061), Seq ID No 31, and deduced translation product as described in Gray, G et al. (1986) J. Bacter **166**:635-643.

Fig.2 shows the amino acid sequence of the mature alpha amylase enzyme from *B. licheniformis* (NCIB8061), Seq ID No 32.

Fig.3 shows an alignment of primary structures of *Bacillus* alpha-amylases. The *B. licheniformis* amylase (Am-Lich), Seq ID No 33, is described by Gray, G. et al. (1986) J.Bact. **166**:635-643, the *B. amyloliquefaciens* amylase (Am-Amylo), Seq ID No 34, is described by Takkinen, K. et al. (1983) J. Biol. Chem. **285**: 1007-1013; and the *B. stearothermophilus* (Am-Stearo), Seq ID No 35, is described by Ihara, H. et al. (1985) J. Biochem, **98**:95-103.

Fig. 4 shows the amino acid sequence of the mature alpha-amylase variant M197T, Seq ID No 36.

Fig. 4b shows the amino acid sequence of the A4 form of alpha-amylase from *B. licheniformis* NCIB8061, Seq ID No 37. Numbering is from the N-terminus, starting with the four additional alanines.

Fig. 5 shows plasmid pA4BL wherein BLAA refers to *B. licheniformis* alpha-amylase gene, PstI to SstI; Amp^R refers to the ampicillin-resistant gene from pBR322; and CAT refers to the Chloramphenicol-resistant gene from pC194.

Fig. 6 shows the signal sequence-mature protein junctions for *B. licheniformis* (Seq ID No 38), *B. subtilis* (Seq ID No 39), *B. licheniformis* in pA4BL (Seq ID No 40) and *B. licheniformis* in pBLapr (Seq ID No 41).

Fig 7 shows inactivation of certain alpha-amylases (Spezyme ® AA20, M15L) with 0.88M H₂O₂ at pH 5.0 25°C.

Fig. 8 shows a schematic for the production of M15X cassette mutants.

Fig. 9 shows expression of M15X variants.

Fig. 10 shows specific activity of M15X variants on soluble starch.

Fig. 11 shows heat stability of M15X variants at 90°C, pH 5.0, 5mM CaCl₂, 5 mins.

Fig. 12 shows a specific activity on starch and soluble substrate, and performance in jet liquefaction at pH 5.5, of M15 variants as a function of percent activity of *B. licheniformis* wild-type.

Fig. 13 shows the inactivation of *B. licheniformis* alpha-amylase (AA20 at 0.65mg/ml) with chloramine-T at pH 8.0 as compared to variants M197A (1.7mg/ml) and M197L (1.7 mg/ml).

Fig. 14 shows the inactivation of *B.licheniformis* alpha-amylase (AA20 at 0.22mg/ml) with chloramine-T at pH 4.0 as compared to variants M197A (4.3 mg/ml) and M197L

Fig. 15 shows the reaction of *B. licheniformis* alpha-amylase (AA20 at 0.75 mg/ml) with chloramine-T at pH 5.0 as compared to double variants M197T/W138F (0.64 mg/ml) and M197T/W138Y (0.60 mg/ml).

It is believed that amylases used in starch liquefaction may be subject to some form of inactivation due to some activity present in the starch slurry (see US applications 07/785,624 and 07/785,623 and US Patent 5,180,669, issued January 19, 1993. Furthermore, use of an amylase in the presence of oxidants, such as in bleach or peracid containing detergents, may result in partial or complete inactivation of the amylase. Therefore, the present invention focuses on altering the oxidative sensitivity of amylases. The mutant enzymes of the present invention may also have an altered pH profile and/or altered thermal stability which may be due to the enhanced oxidative stability of the enzyme at low or high pH's.

Alpha-amylase as used herein includes naturally occurring amylases as well as recombinant amylases. Preferred amylases in the present invention are alpha-amylases derived from *B. licheniformis* or *B. stearothermophilus*, including the A4 form of alpha-amylase derived from *B. licheniformis* as described herein, as well as fungal alpha-amylases as

those derived from *Aspergillus* (i.e. as *A. oryzae* and *A. niger*).

Recombinant alpha-amylases refers to an alpha-amylase in which the DNA sequence encoding the naturally occurring alpha-amylase is modified to produce a mutant DNA sequence which encodes the substitution, insertion or deletion of one or more amino acids in the alpha-amylase sequence. Suitable modification methods are disclosed herein, and also in US Patents 4,760,025 and 5,185,258.

Homologies have been found between almost all endo-amylases sequenced to date, ranging from plants, mammals, and bacteria (Nakajima, R.T. et al. (1986) Appl. Microbiol. Biotechnol. 23:355-360; Rogers, J.C. (1985) Biochem. Biophys. Res. Commun. 128:470-476). There are four areas of particularly high homology in certain *Bacillus* amylases, as shown in Fig. 3, wherein the underlined sections designate the areas of high homology. Further, sequence alignments have been used to map the relationship between *Bacillus* endo-amylases (Feng, D.F. and Doolittle, R.F. (1987) J. Molec. Evol. 35:351-360). The relative sequence homology between *B. stearothermophilus* and *B. licheniformis* amylase is about 66%, as determined by Holm, L. et al. (1990) Protein Engineering 3 (3) pp. 181-191. The sequence homology between *B. licheniformis* and *B. amyloliquefaciens* amylases is about 81%, as per Holm, L. et al., *supra*. While sequence homology is important, it is generally recognized that structural homology is also important in comparing amylases or other enzymes. For example, structural homology between fungal amylases and bacterial (*Bacillus*) amylase have been suggested and, therefore, fungal amylases are encompassed within the present invention.

An alpha-amylase mutant has an amino acid sequence which is derived from the amino acid sequence of a precursor alpha-amylase. The precursor alpha-amylases include naturally occurring alpha-amylases and recombinant alpha-amylases (as defined). The amino acid sequence of the alpha-amylase mutant is derived from the precursor alpha-amylase amino acid sequence by the substitution, deletion or insertion of one or more amino acids of the precursor amino acid sequence. Such modification is of the precursor DNA sequence which encodes the amino acid sequence of the precursor alpha-amylase rather than manipulation of the precursor alpha-amylase enzyme *per se*. Suitable methods for such manipulation of the precursor DNA sequence include methods disclosed herein and in US patent 4,760,025 and 5,185,258

Specific residues corresponding to positions M15 and W138 of *Bacillus licheniformis* alpha-amylase are identified herein for substitution or deletion, as are all methionine, histidine, tryptophan, cysteine and tyrosine positions. The amino acid position number (i.e., +197) refers to the number assigned to the mature *Bacillus licheniformis* alpha-amylase sequence presented in Fig. 2. The invention, however, is not limited to the mutation of this particular mature alpha-amylase (*B. licheniformis*) but extends to precursor alpha-amylases containing amino acid residues at positions which are equivalent to the particular identified residue in *B. licheniformis* alpha-amylase. A residue (amino acid) of a precursor alpha-amylase is equivalent to a residue of *B. licheniformis* alpha-amylase if it is either homologous (i.e., corresponding in position in either primary or tertiary structure) or analogous to a specific residue or portion of that residue in *B. licheniformis* alpha-amylase (i.e., having the same or similar functional capacity to combine, react, or interact chemically or structurally).

In order to establish homology to primary structure, the amino acid sequence of a precursor alpha-amylase is directly compared to the *B. licheniformis* alpha-amylase primary sequence and particularly to a set of residues known to be invariant to all alpha-amylases for which sequence is known, as seen in Fig. 3. It is possible also to determine equivalent residues by tertiary structure: crystal structures have been reported for porcine pancreatic alpha-amylase (Buisson, G. et al. (1987) EMBO J. 6:3909-3916); Taka-amylase A from *Aspergillus oryzae* (Matsuura, Y. et al. (1984) J. Biochem. (Tokyo) 95:697-702); and an acid alpha-amylase from *A. niger* (Boel, E. et al. (1990) Biochemistry 29: 6244-6249), with the former two structures being similar. There are no published structures for *Bacillus* alpha-amylases, although there are predicted to be common super-secondary structures between glucanases (MacGregor, E.A. & Svensson, B. (1989) Biochem. J. 259:145-152) and a structure for the *B. stearothermophilus* enzyme has been modeled on that of Taka-amylase A (Holm, L. et al. (1990) Protein Engineering 3:181-191). The four highly conserved regions shown in Fig. 3 contain many residues thought to be part of the active-site (Matsuura, Y. et al. (1984) J. Biochem. (Tokyo) 95:697-702; Buisson, G. et al. (1987) EMBO J. 6:3909-3916; Vihinen, M. et al. (1990) J. Biochem. 107:267-272) including, in the *licheniformis* numbering, His105; Arg229; Asp231; His235; Glu261 and Asp328.

Expression vector as used herein refers to a DNA construct containing a DNA sequence which is operably linked to a suitable control sequence capable of effecting the expression of said DNA in a suitable host. Such control sequences may include a promoter to effect transcription, an optional operator sequence to control such transcription, a sequence encoding suitable mRNA ribosome-binding sites, and sequences which control termination of transcription and translation. A preferred promoter is the *B. subtilis aprE* promoter. The vector may be a plasmid, a phage particle, or simply a potential genomic insert. Once transformed into a suitable host, the vector may replicate and function independently of the host genome, or may, in some instances, integrate into the genome itself. In the present specification, plasmid and vector are sometimes used interchangeably as the plasmid is the most commonly used form of vector at present. However, the invention is intended to include such other forms of expression vectors which serve equivalent functions and which are, or become, known in the art.

Host strains (or cells) useful in the present invention generally are procaryotic or eucaryotic hosts and include any

transformable microorganism in which the expression of alpha-amylase can be achieved. Specifically, host strains of the same species or genus from which the alpha-amylase is derived are suitable, such as a *Bacillus* strain. Preferably an alpha-amylase negative *Bacillus* strain (genes deleted) and/or an alpha-amylase and protease deleted *Bacillus* strain such as *Bacillus subtilis* strain BG2473 ($\Delta amyE, \Delta apr, \Delta npr$) is used. Host cells are transformed or transfected with vectors constructed using recombinant DNA techniques. Such transformed host cells are capable of either replicating vectors encoding the alpha-amylase and its variants (mutants) or expressing the desired alpha-amylase.

Preferably the mutants of the present invention are secreted into the culture medium during fermentation. Any suitable signal sequence, such as the *aprE* signal peptide, can be used to achieve secretion.

Many of the alpha-amylase mutants of the present invention are useful in formulating various detergent compositions, particularly certain dish care cleaning compositions, especially those cleaning compositions containing known oxidants. Alpha-amylase mutants of the invention can be formulated into known powdered, liquid or gel detergents having pH between 6.5 to 12.0. Suitable granular composition may be made as described in commonly owned US patent applications 07/4290,881, 07/533,721 and 07/957,973. These detergent cleaning compositions can also contain other enzymes, such as known proteases, lipases, cellulases, endoglycosidases or other amylases, as well as builders, stabilizers or other excipients known to those skilled in the art. These enzymes can be present as co-granules or as blended mixes or in any other manner known to those skilled in the art. Furthermore, it is contemplated by the present invention that multiple mutants may be useful in cleaning or other applications. For example, a mutant enzyme having changes at both +15 and +197 may exhibit enhanced performance useful in a cleaning product.

As described previously, alpha-amylase mutants of the present invention may also be useful in the liquefaction of starch. Starch liquefaction, particularly granular starch slurry liquefaction, is typically carried out at near neutral pH's and high temperatures. As described in US applications 07/788,624 and 07/785,623 and US Patent 5,180,669, it appears that an oxidizing agent or inactivating agent of some sort is also present in typical liquefaction processes, which may affect the enzyme activity; thus, in these related patent applications an antioxidant is added to the process to protect the enzyme.

Based on the conditions of a preferred liquefaction process, as described in US applications 07/788,624 and 07/785,623 and US Patent 5,180,669, namely low pH, high temperature and potential oxidation conditions, preferred mutants of the present invention for use in liquefaction processes comprise mutants exhibiting altered pH performance profiles (i.e., low pH profile, pH <6 and preferably pH <5.5), and/or altered thermal stability (i.e., high temperature, about 90°-110°C), and/or altered oxidative stability (i.e., enhanced oxidative stability).

Thus, an improved method for liquefying starch is taught by the present invention, the method comprising liquefying a granular starch slurry from either a wet or dry milling process at a pH from about 4 to 6 by adding an effective amount of an alpha-amylase mutant of the present invention to the starch slurry; optionally adding an effective amount of an antioxidant or other excipient to the slurry; and reacting the slurry for an appropriate time and temperature to liquefy the starch.

The following is presented by way of example and is not to be construed as a limitation to the scope of the claims. Abbreviations used herein, particularly three letter or one letter notations for amino acids are described in Dale, J.W., Molecular Genetics of Bacteria, John Wiley & Sons, (1989) Appendix B.

Experimental

Example 1

Substitutions for the Methionine Residues in *B. licheniformis* Alpha-Amylase

The alpha-amylase gene (Fig. 1) was cloned from *B. licheniformis* NCIB8061 obtained from the National Collection of Industrial Bacteria, Aberdeen, Scotland (Gray, G. et al. (1986) J. Bacteriology **166**:635-643). The 1.72kb PstI-SstI fragment, encoding the last three residues of the signal sequence; the entire mature protein and the terminator region was subcloned into M13MP18. A synthetic terminator was added between the BclI and SstI sites using a synthetic oligonucleotide cassette of the form:

BclI	SstI
5' GATCAAAACATAAAAAACCGGCCTTGGCCCCGCCGGTTTTTTATTATTTTGGAGCT	3'
3' TTTTGTATTTTTTGGCCGGAACCGGGGCGGCCAAAAAATAATAAAAC	5'

Seq ID No 1

designed to contain the *B. amyloliquefaciens* subtilisin transcriptional terminator (Wills et al. (1983) Nucleic Acid Research 11:7911-7925).

Site-directed mutagenesis by oligonucleotides used essentially the protocol of Zoller, M. et al. (1983) Meth. Enzymol. 100:468-500: briefly, 5'-phosphorylated oligonucleotide primers were used to introduce the desired mutations on the M13 single-stranded DNA template using the oligonucleotides listed in Table I to substitute for each of the seven methionines found in *B. licheniformis* alpha-amylase. Each mutagenic oligonucleotide also introduced a restriction endonuclease site to use as a screen for the linked mutation.

TABLE I

Mutagenic Oligonucleotides for the Substitution of the Methionine Residues in *B. licheniformis* Alpha-Amylase

15	5'-T GGG ACG CTG GCG <u>CAG TAC</u> TTT GAA TGG T-3'	MSA ScaI+	Seq ID No 2
20	5'-TG ATG <u>CAG TAC</u> TTT GAA TGG <u>TAC</u> CTG CCC AAT GA-3'	M15L ScaI+ KpnI+	Seq ID No 3
	5'-GAT TAT TTG TTG TAT GCC <u>GAT ATC</u> GAC TAT GAC CAT-3'	M197L EcoRV+	Seq ID No 4
25	5'-CG GGG AAG <u>GAG GCC</u> TTT ACG GTA GCT-3'	M256A SmaI+	Seq ID No 5
30	5'-GC GGC TAT GAC <u>TTA</u> AGG AAA TTG C-3'	M304L AclII+	Seq ID No 6
	5'-C TAC GGG <u>GAT GCA</u> TAC GGG ACG A-3'	M366A NsiI+	Seq ID No 7
35	5'-C TAC GGG GAT TAC TAC GGG <u>ACC</u> AAG GGA GAC TCC C-3'	M365Y SlyI+	Seq ID No 8
40	5'-CC GGT GGG <u>GCC AAG</u> CGG GCC TAT GTT GGC CGG CAA A-3'	M438A SfiI+	Seq ID No 9

Bold letter indicate base changes introduced by oligonucleotide.

Codon changes indicated in the form M8A, where methionine (M) at position +8 has been changed to alanine (A).

Underlining indicates restriction endonuclease site introduced by oligonucleotide.

The heteroduplex was used to transfect *E. coli* mutL cells (Kramer et al. (1984) Cell 38:879) and, after plaque-purification, clones were analyzed by restriction analysis of the RF1's. Positives were confirmed by dideoxy sequencing (Sanger et al. (1977) Proc. Natl. Acad. Sci. U.S.A. 74:5463-5467) and the PstI-SstI fragments for each subcloned into an *E. coli* vector, plasmid pA4BL.

Plasmid pA4BL

Following the methods described in US application 860,468 (Power et al.), a silent PstI site was introduced at
 5 condon + 1 (the first amino-acid following the signal cleavage site) of the *aprE* gene from pS168-1 (Stahl, M.L. and
 Ferrari, E. (1984) J. Bacter. **158**:411-418). The *aprE* promoter and signal peptide region was then cloned out of a
 pJH101 plasmid (Ferrari, F.A. et al. (1983) J. Bacter. **154**:1513-1515) as a HindIII-PstI fragment and subcloned into
 the pUC18-derived plasmid JM102 (Ferrari, E. and Hoch, J.A. (1989) Bacillus, ed. C.R. Harwood, Plenum Pub., pp.
 57-72). Addition of the PstI-SstI fragment from *B. licheniformis* alpha-amylase gave pA4BL (Fig. 5) having the resulting
 10 *aprE* signal peptide-amylase junction as shown in Fig. 6.

Transformation Into *B. subtilis*

pA4BL is a plasmid able to replicate in *E. coli* and integrate into the *B. subtilis* chromosome. Plasmids containing
 different variants were transformed into *B. subtilis* (Anagnostopoulos, C. and Spizizen, J. (1961) J. Bacter. **81**:741-746)
 15 and integrated into the chromosome at the *aprE* locus by a Campbell-type mechanism (Young, M. (1984) J. Gen.
 Microbiol. **130**:1613-1621). The *Bacillus subtilis* strain BG2473 was a derivative of 1168 which had been deleted for
 amylase ($\Delta amyE$) and two proteases (Δapr , Δnpr) (Stahl, M.L. and Ferrari, E., J. Bacter. **158**:411-418 and US Patent
 5,264,366, incorporated herein by reference). After transformation the *sacU32*(Hy) (Henner, D.J. et al. (1988) J. Bacter.
 170:296-300) mutation was introduced by PBS-1 mediated transduction (Hoch, J.A. (1983) **154**:1513-1515).

20 N-terminal analysis of the amylase expressed from pA4BL in *B. subtilis* showed it to be processed having four
 extra alanines at the N-terminus of the secreted amylase protein ("A4 form"). These extra residues had no significant,
 deleterious effect on the activity or thermal stability of the A4 form and in some applications may enhance performance.
 In subsequent experiments the correctly processed forms of the *licheniformis* amylase and the variant M197T were
 made from a very similar construction (see Fig. 6). Specifically, the 5' end of the A4 construction was subcloned on an
 25 EcoRI-SstII fragment, from pA4BL (Fig. 5) into M13BM20 (Boehringer Mannheim) in order to obtain a coding-strand
 template for the mutagenic oligonucleotide below:

30 5' -CAT CAG CGT CCC ATT AAG ATT TGC AGC CTG CGC AGA CAT GTT
 GCT-3'

Seq ID No 10

35 This primer eliminated the codons for the extra four N-terminal alanines, correct forms being screened for by the
 absence of the PstI site. Subcloning the EcoRI-SstII fragment back into the pA4BL vector (Fig. 5) gave plasmid pBLapr.
 The M197T substitution could then be moved, on a SstII-SstI fragment, out of pA4BL (M197T) into the complementary
 pBLapr vector to give plasmid pBLapr (M197T). N-terminal analysis of the amylase expressed from pBLapr in *B. subtilis*
 40 showed it to be processed with the same N-terminus found in *B. licheniformis* alpha-amylase.

Example 2Oxidative Sensitivity of Methionine Variants

45 *B. licheniformis* alpha-amylase, such as Spezyme® AA20 (commercially available from Genencor International,
 Inc.), is inactivated rapidly in the presence of hydrogen peroxide (Fig. 7). Various methionine variants were expressed
 in shake-flask cultures of *B. subtilis* and the crude supernatants purified by ammonium sulphate cuts. The amylase
 was precipitated from a 20% saturated ammonium sulphate supernatant by raising the ammonium sulphate to 70%
 50 saturated, and then resuspended. The variants were then exposed to 0.88M hydrogen peroxide at pH 5.0, at 25°C.
 Variants at six of the methionine positions in *B. licheniformis* alpha-amylase were still subject to oxidation by peroxide
 while the substitution at position +197 (M197L) showed resistance to peroxide oxidation. (See Fig. 7.) However, sub-
 sequent analysis described in further detail below showed that while a variant may be susceptible to oxidation at pH
 5.0, 25°C, it may exhibit altered/enhanced properties under different conditions (i.e., liquefaction).
 55

Example 3

Construction of All Possible Variants at Position 197

5 All of the M197 variants (M197X) were produced in the A4 form by cassette mutagenesis, as outlined in Fig. 8:

1) Site directed mutagenesis (via primer extension in M13) was used to make M197A using the mutagenic oligonucleotide below:

10

M197A
5'-GAT TAT TTG GCG TAT GCC GAT ATC GAC TAT GAC CAT-3'
EcoRV+
ClaI- Seq ID No 11

15

which also inserted an EcoRV site (codons 200-201) to replace the ClaI site (codons 201-202). (codons 201-202).

20

2) Then primer LAAM12 (Table II) was used to introduce another silent restriction site (BstBI) over codons 186-188.

3) The resultant M197A (BstBI +, EcoRV +) variant was then subcloned (PstI-SstI fragment) into plasmid pA4BL and the resultant plasmid digested with BstBI and EcoRV and the large vector-containing fragment isolated by electroelution from agarose gel.

25

4) Synthetic primers LAAM14-30 (Table II) were each annealed with the largely complementary common primer LAAM13 (Table II). The resulting cassettes encoded for all the remaining naturally occurring amino acids at position +197 and were ligated, individually, into the vector fragment prepared above.

30

TABLE II

Synthetic Oligonucleotides Used for Cassette Mutagenesis to Produce M197X Variants

35

LAAM12 GG GAA GTT TCG AAT GAA AAC G Seq ID No 12

LAAM13 X197bs Seq ID No 13
(EcoRV) GTC GGC ATA TG CAT ATA ATC ATA GTT GCC GTT TTC ATT (8st8l)

40

LAAM14 1197 Seq ID No 14
(BslBI) CG AAT GAA AAC GGC AAC TAT GAT TAT TTG ATC TAT GCC GAC (EcoRV-)

LAAM15 F197 Seq ID No 15
(BstBI) CG AAT GAA AAC GGC AAC TAT GAT TAT TTG TTC TAT GCC GAC (EcoRV-)

45

LAAM16 V197 Seq ID No 16
(BslBI) CG AAT GAA AAC GGC AAC TAT GAT TAT TTG GTT TAT GCC GAC (EcoRV-)

LAAM17 S197 Seq ID No 17
(BstBI) CG AAT GAA AAC GGC AAC TAT GAT TAT TTG AGC TAT GCC GAC (EcoRV-)

50

LAAM18 P197 Seq ID No 18
(BstBI) CG AAT GAA AAC GGC AAC TAT GAT TAT TTG CCT TAT GCC GAC (EcoRV-)

LAAM19 T197 Seq ID No 19
(BstBI) CG AAT GAA AAC GGC AAC TAT GAT TAT TTG ACA TAT GCC GAC (EcoRV)

55

LAAM20 Y197 Seq ID No 20
(BslBI) CG AAT GAA AAC GGC AAC TAT GAT TAT TTG TAC TAT GCC GAC (EcoRV)

	AM21	H197 (BstBI) CG AAT GAA AAC GGC AAC TAT GAT TAT TTG <u>CAC</u> TAT GCC GAC (EcoRV-)	Seq ID No 21
5	AM22	G197 (BstBI) CG AAT GAA AAC GGC AAC TAT GAT TAT TTG <u>GGC</u> TAT GCC GAC (EcoRV-)	Seq ID No 22
	AM23	Q197 (BstBI) CG AAT GAA AAC GGC AAC TAT GAT TAT TTG <u>CAA</u> TAT GCC GAC (EcoRV-)	Seq ID No 23
10	AM24	N197 (BstBI) CG AAT GAA AAC GGC AAC TAT GAT TAT TTG <u>AAC</u> TAT GCC GAC (EcoRV-)	Seq ID No 24
	AM25	K197 (BstBI) CG AAT GAA AAC GGC AAC TAT GAT TAT TTG <u>AAA</u> TAT GCC GAC (EcoRV-)	Seq ID No 25
15	AM26	D197 (BstBI) CG AAT GAA AAC GGC AAC TAT GAT TAT TTG <u>GAT</u> TAT GCC GAC (EcoRV-)	Seq ID No 26
	AM27	E197 (BstBI) CG AAT GAA AAC GGC AAC TAT GAT TAT TTG <u>GAA</u> TAT GCC GAC (EcoRV-)	Seq ID No 27
20	AM28	C197 (BstBI) CG AAT GAA AAC GGC AAC TAT GAT TAT TTG <u>TGT</u> TAT GCC GAC (EcoRV-)	Seq ID No 28
	AM29	W197 (BstBI) CG AAT GAA AAC GGC AAC TAT GAT TAT TTG <u>TGG</u> TAT GCC GAC (EcoRV-)	Seq ID No 29
25	AM30	R197 (BstBI) CG AAT GAA AAC GGC AAC TAT GAT TAT TTG <u>AGA</u> TAT GCC GAC (EcoRV-)	Seq ID No 30

The cassettes were designed to destroy the EcoRV site upon ligation, thus plasmids from *E. coli* transformants were screened for loss of this unique site. In addition, the common bottom strand of the cassette contained a frame-shift and encoded a NsiI site, thus transformants derived from this strand could be eliminated by screening for the presence of the unique NsiI site and would not be expected, in any case, to lead to expression of active amylase.

Positives by restriction analysis were confirmed by sequencing and transformed in *B. subtilis* for expression in shake-flask cultures. The specific activity of certain of the M197X mutants was then determined using a soluble substrate assay. The data generated using the following assay methods are presented below in Table III.

Soluble Substrate Assay:

A rate assay was developed based on an end-point assay kit supplied by Megazyme (Aust.) Pty. Ltd.: Each vial of substrate (p-nitrophenyl maltoheptaoside, BPNPG7) was dissolved in 10ml of sterile water, followed by a 1 to 4 dilution in assay buffer (50mM maleate buffer, pH 6.7, 5mM calcium chloride, 0.002% Tween20). Assays were performed by adding 10μl of amylase to 790μl of the substrate in a cuvette at 25°C. Rates of hydrolysis were measured as the rate of change of absorbance at 410nm, after a delay of 75 seconds. The assay was linear up to rates of 0.4 absorption units/min.

The amylase protein concentration was measured using the standard Bio-Rad assay (Bio-Rad Laboratories) based on the method of Bradford, M. (1976) Anal. Biochem. 72:248) using bovine serum albumin standards.

Starch Hydrolysis Assay:

The standard method for assaying the alpha-amylase activity of Spezyme® AA20 was used. This method is described in detail in Example 1 of USSN 07/785,624, incorporated herein by reference. Native starch forms a blue color with iodine but fails to do so when it is hydrolyzed into shorter dextrin molecules. The substrate is soluble Lintner starch 5gm/liter in phosphate buffer, pH 6.2 (42.5gm/liter potassium dihydrogen phosphate, 3.16gm/liter sodium hydroxide). The sample is added in 25mM calcium chloride and activity is measured as the time taken to give a negative iodine test upon incubation at 30°C. Activity is recorded in liquefons per gram or ml (LU) calculated according to the formula:

$$\text{LU/ml or LU/g} = \frac{570}{V \times t} \times D$$

Where

LU = liquefon unit

V = volume of sample (5ml)

t = dextrinization time (minutes)

D = dilution factor = dilution volume/ml or g of added enzyme.

TABLE III

ALPHA-AMYLASE	SPECIFIC ACTIVITY (as % of AA20 value) on:	
	Soluble Substrate	Starch
Spezyme® AA20	100	100
A4 form	105	115
M15L (A4 form)	93	94
M15L	85	103
M197T (A4 form)	75	83
M197T	62	81
M197A (A4 form)	88	89
M197C	85	85
M197L (A4 form)	51	17

Example 4

Characterization of Variant M15L

Variant M15L made as per the prior examples did not show increased amylase activity (Table III) and was still inactivated by hydrogen peroxide (Fig. 7). It did, however, show significantly increased performance in jet-liquefaction of starch, especially at low pH as shown in Table IV below.

Starch liquefaction was typically performed using a Hydroheater M 103-M steam jet equipped with a 2.5 liter delay coil behind the mixing chamber and a terminal back pressure valve. Starch was fed to the jet by a Moyno pump and steam was supplied by a 150 psi steam line, reduced to 90-100 psi. Temperature probes were installed just after the Hydroheater jet and just before the back pressure valve.

Starch slurry was obtained from a corn wet miller and used within two days. The starch was diluted to the desired solids level with deionized water and the pH of the starch was adjusted with 2% NaOH or saturated Na₂CO₃. Typical liquefaction conditions were:

Starch	32%-35% solids
Calcium	40-50 ppm (30 ppm added)
pH	5.0-6.0
Alpha-amylase	12-14 LU/g starch dry basis

Starch was introduced into the jet at about 350 ml/min. The jet temperature was held at 105°-107°C. Samples of starch were transferred from the jet cooker to a 95°C second stage liquefaction and held for 90 minutes.

The degree of starch liquefaction was measured immediately after the second stage liquefaction by determining the dextrose equivalence (DE) of the sample and by testing for the presence of raw starch, both according to the methods described in the Standard Analytical Methods of the Member Companies of the Corn Refiners Association, Inc., sixth edition. Starch, when treated generally under the conditions given above and at pH 6.0, will yield a liquefied starch with a DE of about 10 and with no raw starch. Results of starch liquefaction tests using mutants of the present invention are provided in Table IV.

TABLE IV

Performance of Variants M15L (A4 form) and M15L in Starch Liquefaction		
	pH	DE after 90 Mins.
Spezyme® AA20	5.9	9.9
M15L (A4 form)	5.9	10.4
Spezyme® AA20	5.2	1.2
M15L (A4 form)	5.2	2.2
Spezyme® AA20	5.9	9.3*
M15L	5.9	11.3*
Spezyme® AA20	5.5	3.25**
M15L	5.5	6.7**
Spezyme® AA20	5.2	0.7**
M15L	5.2	3.65**

*average of three experiments

** average of two experiments

Example 5Construction of M15X Variants

Following generally the processes described in Example 3 above, all variants at M15 (M15X) were produced in native *B. licheniformis* by cassette mutagenesis, as outlined in Fig. 8.

1) Site directed mutagenesis (via primer extension in M13) was used to introduce unique restriction sites flanking the M15 codon to facilitate insertion of a mutagenesis cassette. Specifically, a BstB1 site at codons 11-13 and a Msc1 site at codons 18-20 were introduced using the two oligonucleotides shown below.

M15XBstB1 5'-G ATG CAG TAT TTC GAA CTGG TAT A-3'
BstB1

Seq ID No 48

M15XMsc1 5'-TG CCC AAT GAT GGC CAA CAT TGG AAG-3'
Msc1

Seq ID No 49

2) The vector for M15X cassette mutagenesis was then constructed by subcloning the Sfi1-SstII fragment from the mutagenized amylase (BstB1 +, Msc1+) into plasmid pBLapr. The resulting plasmid was then digested with BstB1 and Msc1 and the large vector fragment isolated by electroelution from a polyacrylamide gel.

3) Mutagenesis cassettes were created as with the M197X variants. Synthetic oligomers, each encoding a substitution at codon 15, were annealed to a common bottom primer. Upon proper ligation of the cassette to the vector, the Msc1 is destroyed allowing for screening of positive transformants by loss of this site. The bottom primer contains an unique SnaB1 site allowing for the transformants derived from the bottom strand to be eliminated by screening for the SnaB1 site. This primer also contains a frameshift which would also eliminate amylase expression for the mutants derived from the common bottom strand.

The synthetic cassettes are listed in Table V and the general cassette mutagenesis strategy is illustrated in Figure 8.

TABLE V

Synthetic Oligonucleotides Used for Cassette Mutagenesis
to Produce M15X Variants

5

15A	(BstB1)	C GAA TGG TAT <u>GCT</u> CCC AAT GAC GG (Msc1)	Seq ID No 50
15R	(BstB1)	C GAA TGG TAT <u>CGC</u> CCC AAT GAC GG (Msc1)	Seq ID No 51
10 15N	(BstB1)	C GAA TGG TAT <u>AAT</u> CCC AAT GAC GG (Msc1)	Seq ID No 52
15D	(BstB1)	C GAA TGG TAT <u>GAT</u> CCC AAT GAC GG (Msc1)	Seq ID No 53
15H	(BstB1)	C GAA TGG TAT <u>CAC</u> CCC AAT GAC GG (Msc1)	Seq ID No 54
15 15K	(BstB1)	C GAA TGG TAT <u>AAA</u> CCC AAT GAC GG (Msc1)	Seq ID No 55
15P	(BstB1)	C GAA TGG TAT <u>CCG</u> CCC AAT GAC GG (Msc1)	Seq ID No 56
15S	(BstB1)	C GAA TGG TAT <u>TCT</u> CCC AAT GAC GG (Msc1)	Seq ID No 57
20 15T	(BstB1)	C GAA TGG TAT <u>ACT</u> CCC AAT GAC GG (Msc1)	Seq ID No 58
15V	(BstB1)	C GAA TGG TAT <u>GTT</u> CCC AAT GAC GG (Msc1)	Seq ID No 59
15C	(BstB1)	C GAA TGG TAT <u>TGT</u> CCC AAT GAC GG (Msc1)	Seq ID No 60
25 15Q	(BstB1)	C GAA TGG TAT <u>CAA</u> CCC AAT GAC GG (Msc1)	Seq ID No 61
15E	(BstB1)	C GAA TGG TAT <u>GAA</u> CCC AAT GAC GG (Msc1)	Seq ID No 62
15G	(BstB1)	C GAA TGG TAT <u>GGT</u> CCC AAT GAC GG (Msc1)	Seq ID No 63
30 15I	(BstB1)	C GAA TGG TAT <u>ATT</u> CCC AAT GAC GG (Msc1)	Seq ID No 64
15F	(BstB1)	C GAA TGG TAT <u>TTT</u> CCC AAT GAC GG (Msc1)	Seq ID No 65
15W	(BstB1)	C GAA TGG TAT <u>TGG</u> CCC AAT GAC GG (Msc1)	Seq ID No 66
35 15Y	(BstB1)	C GAA TGG TAT <u>TAT</u> CCC AAT GAC GG (Msc1)	Seq ID No 67
M15X	(Msc1)	CC GTC ATT GGC ACT ACG TAC CAT T (BstB1) (bottom strand)	Seq ID No 68

40

Underline indicates codon changes at amino acid position 15.

Conservative substitutions were made in some cases to prevent introduction of new restriction sites.

45

Example 6

Bench Liquefaction with M15X Variants

50

Eleven alpha-amylase variants with substitutions for M15 made as per Example 5 were assayed for activity, as compared to Spezyme® AA20 (commercially available from Genencor International, Inc.) in liquefaction at pH 5.5 using a bench liquefaction system. The bench scale liquefaction system consisted of a stainless steel coil (0.25 inch diameter, approximately 350 ml volume) equipped with a 7 inch long static mixing element approximately 12 inches from the anterior end and a 30 psi back pressure valve at the posterior end. The coil, except for each end, was immersed

55

in a glycerol-water bath equipped with thermostatically controlled heating elements that maintained the bath at 105-106°C.

Starch slurry containing enzyme, maintained in suspension by stirring, was introduced into the reaction coil by a piston driven metering pump at about 70 ml/min. The starch was recovered from the end of the coil and was transferred

to the secondary hold (95°C for 90 minutes). Immediately after the secondary hold, the DE of the liquefied starch was determined, as described in Example 4. The results are shown in Fig. 12.

Example 7

Characterization of M15X Variants

All M15X variants were propagated in *Bacillus subtilis* and the expression level monitored as shown in Fig. 9. The amylase was isolated and partially purified by a 20-70% ammonium sulfate cut. The specific activity of these variants on the soluble substrate was determined as per Example 3 (Fig. 10). Many of the M15X amylases have specific activities greater than that of Spezyme® AA20. A benchtop heat stability assay was performed on the variants by heating the amylase at 90°C for 5 min. in 50 mM acetate buffer pH 5 in the presence of 5 mM CaCl₂ (Fig. 11). Most of the variants performed as well as Spezyme® AA20 in this assay. Those variants that exhibited reasonable stability in this assay (reasonable stability defined as those that retained at least about 60% of Spezyme® AA20's heat stability) were tested for specific activity on starch and for liquefaction performance at pH 5.5. The most interesting of those mutants are shown in Fig. 16. M15D, N and T, along with L, outperformed Spezyme® AA20 in liquefaction at pH 5.5 and have increased specific activities in both the soluble substrate and starch hydrolysis assays.

Generally, we have found that by substituting for the methionine at position 15, we can provide variants with increased low pH-liquefaction performance and/or increased specific activity.

Example 8

Tryptophan Sensitivity to Oxidation

Chloramine-T (sodium N-chloro-*p*-toluenesulfonimide) is a selective oxidant, which oxidizes methionine to methionine sulfoxide at neutral or alkaline pH. At acidic pH, chloramine-T will modify both methionine and tryptophan (Schechter, Y., Burstein, Y. and Patchornik, A. (1975) *Biochemistry* 14(20) 4497-4503). Fig. 13 shows the inactivation of *B. licheniformis* alpha-amylase with chloramine-T at pH 8.0 (AA20 = 0.65 mg/ml, M197A = 1.7 mg/ml, M197L = 1.7 mg/ml). The data shows that by changing the methionine at position 197 to leucine or alanine, the inactivation of alpha-amylase can be prevented. Conversely, as shown in Fig. 14, at pH 4.0 inactivation of the M197A and M197L proceeds, but require more equivalents of chloramine-T (Fig. 18; AA20 = 0.22 mg/ml, M197A = 4.3 mg/ml, M197L = 0.53 mg/ml; 200 mM NaAcetate at pH 4.0). This suggests that a tryptophan residue is also implicated in the chloramine-T mediated inactivation event. Furthermore, tryptic mapping and subsequent amino acid sequencing indicated that the tryptophan at position 138 was oxidized by chloramine-T (data not shown). To prove this, site-directed mutants were made at tryptophan 138 as provided below:

Preparation of Alpha-Amylase Double Mutants W138 and M197

Certain variants of W138 (F, Y and A) were made as double mutants, with M197T (made as per the disclosure of Example 3). The double mutants were made following the methods described in Examples 1 and 3. Generally, single negative strands of DNA were prepared from an M13MP18 clone of the 1.72kb coding sequence (Pst I-Sst I) of the *B. licheniformis* alpha-amylase M197T mutant. Site-directed mutagenesis was done using the primers listed below, essentially by the method of Zoller, M. et al. (1983) except T4 gene 32 protein and T4 polymerase were substituted for klenow. The primers all contained unique sites, as well as the desired mutation, in order to identify those clones with the appropriate mutation.

Tryptophan 138 to Phenylalanine

133 134 135 136 137 138 139 140 141 142 143
CAC CTA ATT AAA GCT TTC ACA CAT TTT CAT TTT
Hind III

Seq ID No 42

Tryptophan 138 to Tyrosine

133 134 135 136 137 138 139 140 141 142 143
 CAC CTA ATT AAA GCT TAC ACA CAT TTT CAT TTT
 Hind III

Seq ID No 43

Tryptophan 138 to Alanine - This primer also engineers unique sites
 upstream and downstream of the 138 position.

127 128 129 130 131 132 133 134 135 136 137 138 139 140 141 142
 C CGC GTA ATT TCC GGA GAA CAC CTA ATT AAA GCC GCA ACA CAT TTT CAT
 BspE I

143 144 145 146 147
 TTT CCC GGG CGC GGC AG
 Xma I

Seq ID No 44

Mutants were identified by restriction analysis and W138F and W138Y confirmed by DNA sequencing. The W133A sequence revealed a nucleotide deletion between the unique BspE I and Xma I sites, however, the rest of the gene sequenced correctly. The 1.37kb SstII/SstI fragment containing both W138X and M197T mutations was moved from M13MP18 into the expression vector pBLapr resulting in pBLapr (W138F, M197T) and pBLapr (W138Y, M197T). The fragment containing unique BspE I and Xma I sites was cloned into pBLapr (BspE I, Xma I, M197T) since it is useful for cloning cassettes containing other amino acid substitutions at position 138.

Single Mutations at Amino Acid Position 138

Following the general methods described in the prior examples, certain single variants of W138 (F, Y, L, H and C) were made.

The 1.24kb Asp718-SstI fragment containing the M197T mutation in plasmid pBLapr (W138X, M197T) of Example 7 was replaced by the wild-type fragment with methionine at 197, resulting in pBLapr (W138F), pBLapr (W138Y) and pBLapr (BspE I, Xma I).

The mutants W138L, W138H and W138C were made by ligating synthetic cassettes into the pBLapr (BspE I, Xma I) vector using the following primers:

Tryptophan 138 to Leucine

CC GGA GAA CAC CTA ATT AAA GCC CTA ACA CAT TTT CAT TTT C

Seq ID No 45

Tryptophan 138 to Histidine

CC GGA GAA CAC CTA ATT AAA GCC CAC ACA CAT TTT CAT TTT C

Seq ID No 46

Tryptophan 138 to Cysteine

CC GGA GAA CAC CTA ATT AAA GCC TGC ACA CAT TTT CAT TTT C

Seq ID No 47

Reaction of the double mutants M197T/W138F and M197T/W138Y with chloramine-T was compared with wild-type (AA20 = 0.75 mg/ml, M197T/W138F = 0.64 mg/ml, M197T/W138Y = 0.60 mg/ml; 50 mM NaAcetate at pH 5.0). The results shown in Fig. 19 show that mutagenesis of tryptophan 138 has caused the variant to be more resistant to chloramine-T.

Annex to the description

SEQUENCE LISTING

(1) GENERAL INFORMATION:

(i) APPLICANT: GENENCOR INTERNATIONAL, INC.

(ii) TITLE OF INVENTION: Mutant Alpha-Amylase

(iii) NUMBER OF SEQUENCES: 62

(iv) CORRESPONDENCE ADDRESS:

(A) ADDRESSEE: Genencor International, Inc.

(B) STREET: 4 Cambridge Place, 1870 Winton Road South

(C) CITY: Rochester

(D) STATE: NY

(E) COUNTRY: USA

(F) ZIP: 14618

(v) COMPUTER READABLE FORM:

(A) MEDIUM TYPE: Floppy disk

(B) COMPUTER: IBM PC compatible

(C) OPERATING SYSTEM: PC-DOS/MS-DOS

(D) SOFTWARE: PatentIn Release #1.0, Version #1.25

(vi) CURRENT APPLICATION DATA:

(A) APPLICATION NUMBER:

(B) FILING DATE:

(C) CLASSIFICATION:

(vii) ATTORNEY/AGENT INFORMATION:

(A) NAME: Sharon C Baldock

(B) REGISTRATION NUMBER: 3340

(C) REFERENCE/DOCKET NUMBER: 44411/400

(ix) TELECOMMUNICATION INFORMATION:

(A) TELEPHONE: 44 171 404 5921

(B) TELEFAX: 44 171 831 1768

(2) INFORMATION FOR SEQ ID NO:1:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 56 base pairs

(B) TYPE: nucleic acid

(C) STRANDEDNESS: single

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:1:

GATCAAAACA TAAAAAACCG GCCTTGGCCC CGCCGGTTTT TTATTATTTT TGAGCT

56

(2) INFORMATION FOR SEQ ID NO:2:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 19 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

5

(ii) MOLECULE TYPE: DNA (genomic)

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:2:

10

TGGGACGCTG GCGCAGTACT TTGAATGGT

29

(2) INFORMATION FOR SEQ ID NO:3:

(i) SEQUENCE CHARACTERISTICS:

15

(A) LENGTH: 34 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

20

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:3:

TGATGCAGTA CTTTGAATGG TACCTGCCCA ATGA

34

(2) INFORMATION FOR SEQ ID NO:4:

25

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 36 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

30

(ii) MOLECULE TYPE: DNA (genomic)

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:4:

35

GATTATTTGT TGTATGCCGA TATCGACTAT GACCAT

36

(2) INFORMATION FOR SEQ ID NO:5:

(i) SEQUENCE CHARACTERISTICS:

40

(A) LENGTH: 26 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

45

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:5:

CGGGGAAGGA GGCCTTTAGG GTAGCT

26

50

(2) INFORMATION FOR SEQ ID NO:6:

(i) SEQUENCE CHARACTERISTICS:

55

(A) LENGTH: 24 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

5

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:6:

GCGGCTATGA CTTAAGGAAA TTGC

24

10

(2) INFORMATION FOR SEQ ID NO:7:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 13 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

15

(ii) MOLECULE TYPE: DNA (genomic)

20

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:7:

CTACGGGGAT GCATACGGGA CGA

23

(2) INFORMATION FOR SEQ ID NO:8:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 15 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

25

(ii) MOLECULE TYPE: DNA (genomic)

30

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:8:

CTACGGGGAT TACTACGGGA CCAAGGGAGA CTCCC

35

35

(2) INFORMATION FOR SEQ ID NO:9:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 35 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

40

(ii) MOLECULE TYPE: DNA (genomic)

45

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:9:

CCGGTGGGGC CAAGCGGGG TATGTTGGCC GGCAA

36

(2) INFORMATION FOR SEQ ID NO:10:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 45 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

50

(ii) MOLECULE TYPE: DNA (genomic)

55

(x1) SEQUENCE DESCRIPTION: SEQ ID NO:10:
 CATCAGCGTC CCATTAAGAT TTGCAGCCTG CGCAGACATG TTGCT 45

5 (2) INFORMATION FOR SEQ ID NO:11:

(1) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 35 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 10 (D) TOPOLOGY: linear

(11) MOLECULE TYPE: DNA (genomic)

15 (x1) SEQUENCE DESCRIPTION: SEQ ID NO:11:
 GATTATTGG CGTATGCCGA TATCGACTAT GACCAT 36

(2) INFORMATION FOR SEQ ID NO:12:

(1) SEQUENCE CHARACTERISTICS:
 20 (A) LENGTH: 21 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(11) MOLECULE TYPE: DNA (genomic)
 25

(x1) SEQUENCE DESCRIPTION: SEQ ID NO:12:
 GGGAGTTTC GAATGAAGA C 21

30 (2) INFORMATION FOR SEQ ID NO:13:

(1) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 33 base pairs
 (B) TYPE: nucleic acid
 35 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(11) MOLECULE TYPE: DNA (genomic)

40 (x1) SEQUENCE DESCRIPTION: SEQ ID NO:13:
 GTCCGCATAT GCATATAATC ATAGTTGCCG TTTTCATT 33

(2) INFORMATION FOR SEQ ID NO:14:

(1) SEQUENCE CHARACTERISTICS:
 45 (A) LENGTH: 41 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(11) MOLECULE TYPE: DNA (genomic)
 50

(x1) SEQUENCE DESCRIPTION: SEQ ID NO:14:
 55 CGAATGAAAA CGCAGACTAT GATTATTGCA TCTATGCCGA C 41

(2) INFORMATION FOR SEQ ID NO:15:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 41 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:15:

CGAATGAAAA CGGCAACTAT GATTATTTGT TCTATGCCGA C

41

(2) INFORMATION FOR SEQ ID NO:16:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 41 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:16:

CGAATGAAAA CGGCAACTAT GATTATTTGG TTTATGCCGA C

41

(2) INFORMATION FOR SEQ ID NO:17:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 41 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:17:

CGAATGAAAA CGGCAACTAT GATTATTTGA GCTATGCCGA C

41

(2) INFORMATION FOR SEQ ID NO:18:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 41 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:18:

CGAATGAAAA CGGCAACTAT GATTATTTGC CTTATGCCGA C

41

(2) INFORMATION FOR SEQ ID NO:19:

- (i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 41 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:19:

CGAATGAAAA CGGCAACTAT GATTATTTGA CATATGCCGA C 41

(2) INFORMATION FOR SEQ ID NO:20:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 41 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:20:

CGAATGAAAA CGGCAACTAT GATTATTTGT ACTATGCCGA C 41

(2) INFORMATION FOR SEQ ID NO:21:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 41 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:21:

CGAATGAAAA CGGCAACTAT GATTATTTGC ACTATGCCGA C 41

(2) INFORMATION FOR SEQ ID NO:22:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 41 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:22:

CGAATGAAAA CGGCAACTAT GATTATTTGG GCTATGCCGA C 41

(2) INFORMATION FOR SEQ ID NO:23:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 41 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

5 (xi) SEQUENCE DESCRIPTION: SEQ ID NO:23:

CGAATGAAAA CGGCAACTAT GATTATTTGC AATATGCCGA C 41

(2) INFORMATION FOR SEQ ID NO:24:

10 (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 41 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

15 (ii) MOLECULE TYPE: DNA (genomic)

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:24:

20 CGAATGAAAA CGGCAACTAT GATTATTTGA ACTATGCCGA C 41

(2) INFORMATION FOR SEQ ID NO:25:

25 (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 41 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

30 (xi) SEQUENCE DESCRIPTION: SEQ ID NO:25:

GCAATGAAAA CGGCAACTAT GATTATTTGA AATATGCCGA C 41

35 (2) INFORMATION FOR SEQ ID NO:26:

(i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 41 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 40 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

45 (xi) SEQUENCE DESCRIPTION: SEQ ID NO:26:

CGAATGAAAA CGGCAACTAT GATTATTTGG ATTATGCCGA C 41

(2) INFORMATION FOR SEQ ID NO:27:

50 (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 41 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

55 (ii) MOLECULE TYPE: DNA (genomic)

(x1) SEQUENCE DESCRIPTION: SEQ ID NO:27:

CGAATGAAAA CGGCAACTAT GATTATTTGG AATATGCCGA C

41

(2) INFORMATION FOR SEQ ID NO:28:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 41 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

(x1) SEQUENCE DESCRIPTION: SEQ ID NO:28:

CGAATGAAAA CGGCAACTAT GATTATTTGT GTATTGCCGA C

41

(2) INFORMATION FOR SEQ ID NO:29:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 41 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

(x1) SEQUENCE DESCRIPTION: SEQ ID NO:29:

CGAATGAAAA CGGCAACTAT GATTATTTGT GGTATGCCGA C

41

(2) INFORMATION FOR SEQ ID NO:30:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 41 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

(x1) SEQUENCE DESCRIPTION: SEQ ID NO:30:

CGAATGAAAA CGGCAACTAT GATTATTTGA GATATGCCGA C

41

(2) INFORMATION FOR SEQ ID NO:31:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 1968 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

(x1) SEQUENCE DESCRIPTION: SEQ ID NO:31:

AGCTTGAAGA AGTGAAGAAAG CAGAGAGGCT ATTGAATAAA TGAGTAGAAA GCGCCATATC 50

EP 0 867 504 A1

5 GCGGCTTTTC TTTTGGAAGA AAATATAGGG AAAATGGTAC TTGTTAAAAA TTCGGAATAT 120
 TTATACAACA TCATATGTTT CACATTGAAA GGGGAGGAGA ATCATGAAAC AACAAAAACG 180
 10 CCTTTACGCC CGATTGCTGA CGCTGTTATT TCGGCTCATC TTCTTGCTGC CTCATTCTGC 240
 AGCAGCGGCG GCAAATCTTA ATGGGACGCT GATGCAGTAT TTTGAATGGT ACATGCCCAA 300
 TGACGGCCAA CATTGGAAGC GTTTGCAAAA CGACTCGGCA TATTTGGCTG AACACGGTAT 360
 15 TACTGCCGTC TGGATTCCCC CGGCATATAA GGAACCGAGC CAAGCGGATG TGGGCTACGG 420
 TGCTTACGAC CTTTATGATT TAGGGGAGTT TCATCAAAA GGGACGGTTC GGACAAAGTA 480
 CGGCACAAAA GGAGAGCTGC AATCTGCGAT CAAAAGTCTT CATTCCCGCG ACATTAACGT 540
 20 TTACGGGGAT GTGCTCATCA ACCACAAAGG CGGCGCTGAT GCGACCGAAG ATGTAACCGC 600
 GGTGGAAGTC GATCCCGCTG ACCGCAACCG CGTAATTTCA GGAGAACACC TAATTAAAGC 660
 CTGGACACAT TTTCATTTTC CGGGGCGCGG CAGCACATAC ACCGATTTTA AATGGCATTG 720
 GTACCATTTT GACGGAACCG ATTGGGACGA GTCCCGAAAG CTGAACCGCA TCTATAAGTT 780
 TCAAGGAAAG GCTTGGGATT GGAAGTTTC CAATGAAAC GCGAACTATG ATTATTTGAT 840
 GTATGCCGAC ATCGATTATG ACCATCCTGA TGTCGCAGCA GAAATTAAGA GATGGGGCAC 900
 25 TTGGTATGCC AATGAACTGC AATTGGACGG TTTCCGTCTT GATGCTGTCA AACACATTAA 960
 ATTTTCTTTT TTGCGGGATT GGGTTAATCA TGTCAGGGAA AAAACGGCGA AGGAAATGTT 1020
 TACGGTAGCT GAATATTGGC AGAATGACTT GGGCGCGCTG GAAACTATT TGAACAAAC 1080
 AAATTTTAAT CATTCACTGT TTGACGTGCC GCTTCATTAT CAGTTCCATC CTCATCGAC 1140
 30 ACAGGGAGGC GGCTATGATA TGAGGAAATT GCTGAACGGT ACCGTCTTTT CCAAGCATCC 1200
 GTTGAAATCG GTTACATTTG TCGATAACCA TGATACACAG CCGGGGCAAT CGCTTGAGTC 1260
 GACTGTCCAA ACATGGTTTA AGCCGCTTGC TTACGCTTTT ATTCTCACA GGAATCTGC 1320
 35 ATACCCTCAG GTTTTCTACG GGGATATGTA CGGGACGAAA GGAGACTCCC AGCGCGAAAT 1380
 TCCTGCCTTG AAACACAAA TTGAACCGAT CTAAAAGCG AGAAACAGT ATGCGTACGG 1440
 AGCACAGCAT GATTATTTCC ACCACCATGA CATTGTGGG TGGACAAGG AAGCGACAG 1500
 40 CTCGGTTGCA AATTCAGGT TGGCGGCATT AATAACAGAC GGACCCGGTG GGGCAAAGCG 1560
 AATGTATGTC GCGCGGCAAA AGCCCGCTGA GACATGGCAT GACATTACCG GAAACCGTTC 1620
 GGAGCCCGTT CTCATCAATT CGGAAGGCTG GGGAGAGTTT CACGTAAACG GCGGGTCCGT 1680
 45 TTCAATTTAT GTTCAAAGAT AGAAGAGCAG AGAGGACGGA TTCTCTGAAG GAAATCCGTT 1740
 TTTTATTTT GCGCGTCTTA TAAATTTCTT TGATTACATT TTATAATTAA TTTTAACAA 1800
 GTGTCATCAG CCTCAGGAA GGAAGGCTG ACAGTTTCAA TCGCATAGGT AAGCGGGGA 1860
 50 TGAAATGGCA ACGTTATCTG ATGTTCAAA GAAAGCAAAT GTGTCGAAAA TGACGGTATC 1920
 CCGGGTGATC AATCATCCTG AGACTGTGAC GGATGAATTG AAAAAGCT 1968

(2) INFORMATION FOR SEQ ID NO:32:

55 (1) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 493 amino acids

(B) TYPE: amino acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

5

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:32:

10

Ala Asn Leu Asn Gly Thr Leu Met Gln Tyr Phe Glu Trp Tyr Met Pro
 1 5 10 15

Asn Asp Gly Gln His Trp Lys Arg Leu Gln Asn Asp Ser Ala Tyr Leu
 20 25 30

15

Ala Glu His Gly Ile Thr Ala Val Trp Ile Pro Pro Ala Tyr Lys Gly
 35 40 45

Thr Ser Gln Ala Asp Val Gly Tyr Gly Ala Tyr Asp Leu Tyr Asp Leu
 50 55 60

20

Gly Glu Phe His Gln Lys Gly Thr Val Arg Thr Lys Tyr Gly Thr Lys
 65 70 75 80

Gly Glu Leu Gln Ser Ala Ile Lys Ser Leu His Ser Arg Asp Ile Asn
 85 90 95

25

Val Tyr Gly Asp Val Val Ile Asn His Lys Gly Gly Ala Asp Ala Thr
 100 105 110

Glu Asp Val Thr Ala Val Glu Val Asp Pro Ala Asp Arg Asn Arg Val
 115 120 125

30

Ile Ser Gly Glu His Leu Ile Lys Ala Trp Thr His Phe His Phe Pro
 130 135 140

Gly Arg Gly Ser Thr Tyr Ser Asp Phe Lys Trp His Trp Tyr His Phe
 145 150 155 160

35

Asp Gly Thr Asp Trp Asp Glu Ser Arg Lys Leu Asn Arg Ile Tyr Lys
 165 170 175

Phe Gln Gly Lys Ala Trp Asp Trp Glu Val Ser Asn Glu Asn Gly Asn
 180 185 190

40

Tyr Asp Tyr Leu Met Tyr Ala Asp Ile Asp Tyr Asp His Pro Asp Val
 195 200 205

Ala Ala Glu Ile Lys Arg Trp Gly Thr Trp Tyr Ala Asn Glu Leu Gln
 210 215 220

45

Leu Asp Gly Phe Arg Leu Asp Ala Val Lys His Ile Lys Phe Ser Phe
 225 230 235 240

Leu Arg Asp Trp Val Asn His Val Arg Glu Lys Thr Gly Lys Glu Met
 245 250 255

50

Phe Thr Val Ala Glu Tyr Trp Gln Asn Asp Leu Gly Ala Leu Glu Asn
 260 265 270

Tyr Leu Asn Lys Thr Asn Phe Asn His Ser Val Phe Asp Val Pro Leu
 275 280 285

His Tyr Gln Phe His Ala Ala Ser Thr Gln Gly Gly Gly Tyr Asp Met
 290 295 300

55

Arg Lys Leu Leu Asn Gly Thr Val Val Ser Lys His Pro Leu Lys Ser

EP 0 867 504 A1

	305		310		315		320
5	Val Thr Phe Val Asp Asn His Asp Thr Gln Pro Gly Gln Ser Leu Glu	325		330		335	
	Ser Thr Val Gln Thr Trp Phe Lys Pro Leu Ala Tyr Ala Phe Ile Leu	340		345		350	
10	Thr Arg Glu Ser Gly Tyr Pro Gln Val Phe Tyr Gly Asp Met Tyr Gly	355		360		365	
	Thr Lys Gly Asp Ser Gln Arg Glu Ile Pro Ala Leu Lys His Lys Ile	370		375		380	
15	Glu Pro Ile Leu Lys Ala Arg Lys Gln Tyr Ala Tyr Gly Ala Gln His	385		390		395	400
	Asp Tyr Phe Asp His His Asp Ile Val Gly Trp Thr Arg Glu Gly Asp	405		410		415	
20	Ser Ser Val Ala Asn Ser Gly Leu Ala Ala Leu Ile Thr Asp Gly Pro	420		425		430	
	Gly Gly Ala Lys Arg Met Tyr Val Gly Arg Gln Asn Ala Gly Glu Thr	435		440		445	
25	Trp His Asp Ile Thr Gly Asn Arg Ser Glu Pro Val Val Ile Asn Ser	450		455		460	
	Glu Gly Trp Gly Glu Phe His Val Asn Gly Gly Ser Val Ser Ile Tyr	465		470		475	480
	Val Gln Arg						

(2) INFORMATION FOR SEO ID NO: 11:

(1) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 511 amino acids
(B) TYPE: amino acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:33:

Met Lys Gln Gln Lys Arg Leu Tyr Ala Arg Leu Leu Thr Leu Leu Phe
1 5 10 15

Ala Leu Ile Phe Leu Leu Pro His Ser Ala Ala Ala Ala Ala Asn Leu
20 25 30

Asn Gly Thr Leu Met Gln Tyr Phe Glu Trp Tyr Met Pro Asn Asp Gly
33 40 45

His Trp Lys Arg Leu Gln Asn Asp Ser Ala Tyr Leu Ala Glu His Gly
50 55 60

Ile Thr Ala Val Trp Ile Pro Pro Ala Tyr Lys Gly Thr Ser Gln Ala
65 70 75 80

Asp Val Gly Tyr Gly Ala Tyr Asp Leu Tyr Asp Leu Gly Glu Phe His
35 90 95

Gln Lys Gly Thr Val Arg Thr Lys Trp Gly Thr Lys Gly Glu Leu Gln

EP 0 867 504 A1

	100	105	110
5	Ser Ala Ile Lys Ser Leu His 115	Ser Arg Asp Ile Asn 120	Val Tyr Gly Asp 125
	Val Val Ile Asn His Lys Gly 130	Gly Ala Asp Ala Thr 135	Glu Asp Val Thr 140
10	Ala Val Glu Val Asp Pro Ala Asp Arg Asn Arg Val Ile Ser Gly Glu 145	150	155
	His Leu Ile Lys Ala Trp Thr His Phe His Phe Pro Gly Arg Gly Ser 165	170	175
15	Thr Tyr Ser Asp Phe Lys Trp His Trp Tyr His Phe Asp Gly Thr Asp 180	185	190
	Trp Asp Glu Ser Arg Lys Leu Asn Arg Ile Tyr Lys Phe Gln Gly Lys 195	200	205
20	Ala Trp Asp Trp Glu Val Ser Asn Glu Asn Gly Asn Tyr Asp Tyr Leu 210	215	220
	Met Tyr Ala Asp Ile Asp Tyr Asp His Pro Asp Val Ala Ala Glu Ile 225	230	235
25	Lys Arg Trp Gly Thr Trp Tyr Ala Asn Glu Leu Gln Leu Asp Gly Phe 245	250	255
	Arg Leu Asp Ala Val Lys His Ile Lys Phe Ser Phe Leu Arg Asp Trp 260	265	270
30	Val Asn His Val Arg Glu Lys Thr Gly Lys Glu Met Phe Thr Val Ala 275	280	285
	Glu Tyr Trp Gln Asn Asp Leu Gly Ala Leu Glu Asn Tyr Leu Asn Lys 290	295	300
35	Thr Asn Phe Asn His Ser Val Phe Asp Val Pro Leu His Tyr Gln Phe 305	310	315
	His Ala Ala Ser Thr Gln Gly Gly Gly Tyr Asp Met Arg Lys Leu Leu 325	330	335
40	Asn Gly Thr Val Val Ser Lys His Pro Leu Lys Ser Val Thr Phe Val 340	345	350
	Asp Asn His Asp Thr Gln Pro Gly Gln Ser Leu Glu Ser Thr Val Gln 355	360	365
45	Thr Trp Phe Lys Pro Leu Ala Tyr Ala Phe Ile Leu Thr Arg Glu Ser 370	375	380
	Gly Tyr Pro Gln Val Phe Tyr Gly Asp Met Tyr Gly Thr Lys Gly Asp 385	390	395
50	Ser Gln Arg Glu Ile Pro Ala Leu Lys His Lys Ile Glu Pro Ile Leu 405	410	415
	Lys Ala Arg Lys Gln Tyr Ala Tyr Gly Ala Gln His Asp Tyr Phe Asp 420	425	430
55	His His Asp Ile Val Gly Trp Thr Arg Glu Gly Asp Ser Ser Val Ala 435	440	445
	Asn Ser Gly Leu Ala Ala Leu Ile Thr Asp Gly Pro Gly Gly Ala Lys 450	455	460

EP 0 867 504 A1

Arg Met Tyr Val Gly Arg Gln Asn Ala Gly Glu Thr Trp His Asp Ile
465 470 475 480

Thr Gly Asn Arg Ser Glu Pro Val Val Ile Asn Ser Glu Gly Trp Gly
485 490 495

Glu Phe His Val Asn Gly Gly Ser Val Ser Ile Tyr Val Gln Arg
500 505 510

(2) INFORMATION FOR SEQ ID NO:34:

(i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 520 amino acids
(B) TYPE: amino acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:34:

Met Arg Gly Arg Gly Asn Met Ile Gln Lys Arg Lys Arg Thr Val Ser
1 5 10 15

Phe Arg Leu Val Leu Met Cys Thr Leu Leu Phe Val Ser Leu Pro Ile
20 25 30

Thr Lys Thr Ser Ala Val Asn Gly Thr Leu Met Gln Tyr Phe Glu Trp
35 40 45

Tyr Thr Pro Asn Asp Gly Gln His Trp Lys Arg Leu Gln Asn Asp Ala
50 55 60

Glu His Leu Ser Asp Ile Gly Ile Thr Ala Val Trp Ile Pro Pro Ala
65 70 75 80

Tyr Lys Gly Leu Ser Gln Ser Asp Asn Gly Tyr Gly Pro Tyr Asp Leu
85 90 95

Tyr Asp Leu Gly Glu Phe Gln Gln Lys Gly Thr Val Arg Thr Lys Tyr
100 105 110

Gly Thr Lys Ser Glu Leu Gln Asp Ala Ile Gly Ser Leu His Ser Arg
115 120 125

Asn Val Gln Val Tyr Gly Asp Val Val Leu Asn His Lys Ala Gly Ala
130 135 140

Asp Ala Thr Glu Asp Val Thr Ala Val Glu Val Asn Pro Ala Asn Arg
145 150 155 160

Asn Gln Glu Thr Ser Glu Glu Tyr Gln Ile Lys Ala Trp Thr Asp Phe
165 170 175

Arg Phe Pro Gly Arg Gly Asn Thr Tyr Ser Asp Phe Lys Trp His Trp
180 185 190

Tyr His Phe Asp Gly Ala Asp Trp Asp Glu Ser Arg Lys Ile Ser Arg
195 200 205

Ile Phe Lys Phe Arg Gly Glu Gly Lys Ala Trp Asp Trp Glu Val Ser
210 215 220

Ser Glu Asn Gly Asn Tyr Asp Tyr Leu Met Tyr Ala Asp Val Asp Tyr
225 230 235 240

EP 0 867 504 A1

Asp His Pro Asp Val Val Ala Glu Thr Lys Lys Trp Gly Ile Trp Tyr
 245 250 255
 5 Ala Asn Glu Leu Ser Leu Asp Gly Phe Arg Ile Asp Ala Ala Lys His
 250 265 270
 Ile Lys Phe Ser Phe Leu Arg Asp Trp Val Gln Ala Val Arg Gln Ala
 275 280 285
 10 Thr Gly Lys Glu Met Phe Thr Val Ala Glu Tyr Trp Gln Asn Asn Ala
 290 295 300
 Gly Lys Leu Glu Asn Tyr Leu Asn Lys Thr Ser Phe Asn Gln Ser Val
 305 310 315 320
 15 Phe Asp Val Pro Leu His Phe Asn Leu Gln Ala Ala Ser Ser Gln Gly
 325 330 335
 Gly Gly Tyr Asp Met Arg Arg Leu Leu Asp Gly Thr Val Val Ser Arg
 340 345 350
 20 His Pro Glu Lys Ala Val Thr Phe Val Glu Asn His Asp Thr Gln Pro
 355 360 365
 Gly Gln Ser Leu Glu Ser Thr Val Gln Thr Trp Phe Lys Pro Leu Ala
 370 375 380
 25 Tyr Ala Phe Ile Leu Thr Arg Glu Ser Gly Tyr Pro Gln Val Phe Tyr
 385 390 395 400
 Gly Asp Met Tyr Gly Thr Lys Gly Thr Ser Pro Lys Glu Ile Pro Ser
 405 410 415
 30 Leu Lys Asp Asn Ile Glu Pro Ile Leu Lys Ala Arg Lys Glu Tyr Ala
 420 425 430
 Tyr Gly Pro Gln His Asp Tyr Ile Asp His Pro Asp Val Ile Gly Trp
 435 440 445
 Thr Arg Glu Gly Asp Ser Ser Ala Ala Lys Ser Gly Leu Ala Ala Leu
 450 455 460
 35 Ile Thr Asp Gly Pro Gly Gly Ser Lys Arg Met Tyr Ala Gly Leu Lys
 465 470 475 480
 Asn Ala Gly Glu Thr Trp Tyr Asp Ile Thr Gly Asn Arg Ser Asp Thr
 485 490 495
 40 Val Lys Ile Gly Ser Asp Gly Trp Gly Glu Phe His Val Asn Asp Gly
 500 505 510
 Ser Val Ser Ile Tyr Val Gln Lys
 515 520

(2) INFORMATION FOR SEQ ID NO:35:

(i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 543 amino acids
 (B) TYPE: amino acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(x1) SEQUENCE DESCRIPTION: SEQ ID NO:35:

EP 0 867 504 A1

	Val	Leu	Thr	Phe	His	Arg	Ile	Ile	Arg	Lys	Gly	Trp	Met	Phe	Leu	Leu	
	1				5					10					15		
5	Ala	Phe	Leu	Leu	Thr	Ala	Ser	Leu	Phe	Cys	Pro	Thr	Gly	Arg	His	Ala	
				20					25					30			
	Lys	Ala	Ala	Ala	Pro	Phe	Asn	Gly	Thr	Met	Met	Gln	Tyr	Phe	Glu	Trp	
			35					40					45				
10	Tyr	Leu	Pro	Asp	Asp	Gly	Thr	Leu	Trp	Thr	Lys	Val	Ala	Asn	Glu	Ala	
		50					55					60					
	Asn	Asn	Leu	Ser	Ser	Leu	Gly	Ile	Thr	Ala	Leu	Ser	Leu	Pro	Pro	Ala	
	65					70					75				80		
15	Tyr	Lys	Gly	Thr	Ser	Arg	Ser	Asp	Val	Gly	Tyr	Gly	Val	Tyr	Asp	Leu	
					85					90					95		
	Tyr	Asp	Leu	Gly	Glu	Phe	Asn	Gln	Lys	Gly	Thr	Val	Arg	Thr	Lys	Tyr	
				100					105					110			
20	Gly	Thr	Lys	Ala	Gln	Tyr	Leu	Gln	Ala	Ile	Gln	Ala	Ala	His	Ala	Ala	
				115				120					125				
	Gly	Met	Gln	Val	Tyr	Ala	Asp	Val	Val	Phe	Asp	His	Lys	Gly	Gly	Ala	
		130					135					140					
25	Asp	Gly	Thr	Glu	Trp	Val	Asp	Ala	Val	Glu	Val	Asn	Pro	Ser	Asp	Arg	
	145					150					155					160	
	Asn	Gln	Glu	Ile	Ser	Gly	Thr	Tyr	Gln	Ile	Gln	Ala	Trp	Thr	Lys	Phe	
				165					170						175		
30	Asp	Phe	Pro	Gly	Arg	Gly	Asn	Thr	Tyr	Ser	Ser	Phe	Lys	Trp	Arg	Trp	
				180					185					190			
	Tyr	His	Phe	Asp	Gly	Val	Asp	Trp	Asp	Glu	Ser	Arg	Lys	Leu	Ser	Arg	
			195				200						205				
35	Ile	Tyr	Lys	Phe	Arg	Gly	Ile	Gly	Lys	Ala	Trp	Asp	Trp	Glu	Val	Asp	
		210					215					220					
	Thr	Glu	Asn	Gly	Asn	Tyr	Asp	Tyr	Leu	Met	Tyr	Ala	Asp	Leu	Asp	Met	
	225					230					235				240		
40	Asp	His	Pro	Glu	Val	Val	Thr	Glu	Leu	Lys	Asn	Trp	Gly	Lys	Trp	Tyr	
				245						250					255		
	Val	Asn	Thr	Thr	Asn	Ile	Asp	Gly	Phe	Arg	Leu	Asp	Gly	Leu	Lys	His	
				260					265					270			
45	Ile	Lys	Phe	Ser	Phe	Phe	Pro	Asp	Trp	Leu	Ser	Tyr	Val	Arg	Ser	Gln	
			275					280					285				
	Thr	Gly	Lys	Pro	Leu	Phe	Thr	Val	Gly	Glu	Tyr	Trp	Ser	Tyr	Asp	Ile	
		290					295					300					
50	Asn	Lys	Leu	His	Asn	Tyr	Ile	Thr	Lys	Thr	Asn	Gly	Thr	Met	Ser	Leu	
	305					310					315				320		
	Phe	Asp	Ala	Pro	Leu	His	Asn	Lys	Phe	Tyr	Thr	Ala	Ser	Lys	Ser	Gly	
				325					330						335		
55	Gly	Ala	Phe	Asp	Met	Arg	Thr	Leu	Met	Thr	Asn	Thr	Leu	Met	Lys	Asp	
				340					345				350				
	Gln	Pro	Thr	Leu	Ala	Val	Thr	Phe	Val	Asp	Asn	His	Asp	Thr	Asn	Pro	
			355					360					365				

EP 0 867 504 A1

Ala Lys Arg Cys Ser His Gly Arg Pro Trp Phe Lys Pro Leu Ala Tyr
370 375 380

Ala Phe Ile Leu Thr Arg Gln Glu Gly Tyr Pro Cys Val Phe Tyr Gly
385 390 395 400

Asp Tyr Tyr Gly Ile Pro Gln Tyr Asn Ile Pro Ser Leu Lys Ser Lys
405 410 415

Ile Asp Pro Leu Leu Ile Ala Arg Arg Asp Tyr Ala Tyr Gly Thr Gln
420 425 430

His Asp Tyr Leu Asp His Ser Asp Ile Ile Gly Trp Thr Arg Glu Gly
435 440 445

Val Thr Glu Lys Pro Gly Ser Gly Leu Ala Ala Leu Ile Thr Asp Gly
450 455 460

Ala Gly Arg Ser Lys Trp Met Tyr Val Gly Lys Gln His Ala Gly Lys
465 470 475 480

Val Phe Tyr Asp Leu Thr Gly Asn Arg Ser Asp Thr Val Thr Ile Asn
485 490 495

Ser Asp Gly Trp Gly Glu Phe Lys Val Asn Gly Gly Ser Val Ser Val
500 505 510

Trp Val Pro Arg Lys Thr Thr Val Ser Thr Ile Ala Arg Pro Ile Thr
515 520 525

Thr Arg Pro Trp Thr Gly Glu Phe Val Arg Trp His Glu Pro Arg Leu
530 535 540

Val Ala Trp Pro
545

(2) INFORMATION FOR SEQ ID NO:36:

- (i) SEQUENCE CHARACTERISTICS:
- (A) LENGTH: 483 amino acids
 - (B) TYPE: amino acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:36:

Ala Asn Leu Asn Gly Thr Leu Met Gln Tyr Phe Glu Trp Tyr Met Pro
1 5 10 15

Asn Asp Gly Gln His Trp Lys Arg Leu Gln Asn Asp Ser Ala Tyr Leu
20 25 30

Ala Glu His Gly Ile Thr Ala Val Trp Ile Pro Pro Ala Tyr Lys Gly
35 40 45

Thr Ser Gln Ala Asp Val Gly Tyr Gly Ala Tyr Asp Leu Tyr Asp Leu
50 55 60

Gly Glu Phe His Gln Lys Gly Thr Val Arg Thr Lys Tyr Gly Thr Lys
65 70 75 80

Gly Glu Leu Gln Ser Ala Ile Lys Ser Leu His Ser Arg Asp Ile Asn
85 90 95

EP 0 867 504 A1

	Val	Tyr	Gly	Asp	Val	Val	Ile	Asn	His	Lys	Gly	Gly	Ala	Asp	Ala	Thr	
				100					105						110		
5	Glu	Asp	Val	Thr	Ala	Val	Glu	Val	Asp	Pro	Ala	Asp	Arg	Asn	Arg	Val	
			115					120					125				
	Ile	Ser	Gly	Glu	His	Leu	Ile	Lys	Ala	Trp	Thr	His	Phe	His	Phe	Pro	
		130					135					140					
10	Gly	Arg	Gly	Ser	Thr	Tyr	Ser	Asp	Phe	Lys	Trp	His	Trp	Tyr	His	Phe	
	145					150					155					160	
	Asp	Gly	Thr	Asp	Trp	Asp	Glu	Ser	Arg	Lys	Leu	Asn	Arg	Ile	Tyr	Lys	
				165					170						175		
15	Phe	Gln	Gly	Lys	Ala	Trp	Asp	Trp	Glu	Val	Ser	Asn	Glu	Asn	Gly	Asn	
			180						185					190			
	Tyr	Asp	Tyr	Leu	Thr	Tyr	Ala	Asp	Ile	Asp	Tyr	Asp	His	Pro	Asp	Val	
			195					200					205				
20	Ala	Ala	Glu	Ile	Lys	Arg	Trp	Gly	Thr	Trp	Tyr	Ala	Asn	Glu	Leu	Gln	
		210					215					220					
	Leu	Asp	Gly	Phe	Arg	Leu	Asp	Ala	Val	Lys	His	Ile	Lys	Phe	Ser	Phe	
	225					230					235					240	
25	Leu	Arg	Asp	Trp	Val	Asn	His	Val	Arg	Glu	Lys	Thr	Gly	Lys	Glu	Met	
				245					250						255		
	Phe	Thr	Val	Ala	Glu	Tyr	Trp	Gln	Asn	Asp	Leu	Gly	Ala	Leu	Glu	Asn	
			260					265						270			
30	Tyr	Leu	Asn	Lys	Thr	Asn	Phe	Asn	His	Ser	Val	Phe	Asp	Val	Pro	Leu	
		275					280						285				
	His	Tyr	Gln	Phe	His	Ala	Ala	Ser	Thr	Gln	Gly	Gly	Gly	Tyr	Asp	Met	
		290					295					300					
35	Arg	Lys	Leu	Leu	Asn	Gly	Thr	Val	Val	Ser	Lys	His	Pro	Leu	Lys	Ser	
	305				310						315					320	
	Val	Thr	Phe	Val	Asp	Asn	His	Asp	Thr	Gln	Pro	Gly	Gln	Ser	Leu	Glu	
				325					330						335		
40	Ser	Thr	Val	Gln	Thr	Trp	Phe	Lys	Pro	Leu	Ala	Tyr	Ala	Phe	Ile	Leu	
			340					345						350			
	Thr	Arg	Glu	Ser	Gly	Tyr	Pro	Gln	Val	Phe	Tyr	Gly	Asp	Met	Tyr	Gly	
		355					360						365				
45	Thr	Lys	Gly	Asp	Ser	Gln	Arg	Glu	Ile	Pro	Ala	Leu	Lys	His	Lys	Ile	
		370					375					380					
	Glu	Pro	Ile	Leu	Lys	Ala	Arg	Lys	Gln	Tyr	Ala	Tyr	Gly	Ala	Gln	His	
	385				390					395						400	
50	Asp	Tyr	Phe	Asp	His	His	Asp	Ile	Val	Gly	Trp	Thr	Arg	Glu	Gly	Asp	
				405						410					415		
	Ser	Ser	Val	Ala	Asn	Ser	Gly	Leu	Ala	Ala	Leu	Ile	Thr	Asp	Gly	Pro	
			420					425						430			
55	Gly	Gly	Ala	Lys	Arg	Met	Tyr	Val	Gly	Arg	Gln	Asn	Ala	Gly	Glu	Thr	
		435						440					445				
	Trp	His	Asp	Ile	Thr	Gly	Asn	Arg	Ser	Glu	Pro	Val	Val	Ile	Asn	Ser	
	450						455					460					

Glu Gly Trp Gly Glu Phe His Val Asn Gly Gly Ser Val Ser Ile Tyr
465 470 475 480

Val Gln Arg

(2) INFORMATION FOR SEQ ID NO:37:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 437 amino acids
(B) TYPE: amino acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:37:

Ala Ala Ala Ala Ala Asn Leu Asn Gly Thr Leu Met Gln Tyr Phe Glu
1 5 10 15
Trp Tyr Met Pro Asn Asp Gly Gln His Trp Lys Arg Leu Gln Asn Asp
20 25 30
Ser Ala Tyr Leu Ala Glu His Gly Ile Thr Ala Val Trp Ile Pro Pro
35 40 45
Ala Tyr Lys Gly Thr Ser Gln Ala Asp Val Gly Tyr Gly Ala Tyr Asp
50 55 60
Leu Tyr Asp Leu Gly Glu Phe His Gln Lys Gly Thr Val Arg Thr Lys
65 70 75 80
Tyr Gly Thr Lys Gly Glu Leu Gln Ser Ala Ile Lys Ser Leu His Ser
95 100 105 110
Arg Asp Ile Asn Val Tyr Gly Asp Val Val Ile Asn His Lys Gly Gly
115 120 125
Ala Asp Ala Thr Glu Asp Val Thr Ala Val Glu Val Asp Pro Ala Asp
130 135 140
Arg Asn Arg Val Ile Ser Gly Glu His Leu Ile Lys Ala Trp Thr His
145 150 155 160
Trp Tyr His Phe Asp Gly Thr Asp Trp Asp Glu Ser Arg Lys Leu Asn
165 170 175
Arg Ile Tyr Lys Phe Gln Gly Lys Ala Trp Asp Trp Glu Val Ser Asn
180 185 190
Glu Asn Gly Asn Tyr Asp Tyr Leu Met Tyr Ala Asp Ile Asp Tyr Asp
195 200 205
His Pro Asp Val Ala Ala Glu Ile Lys Arg Trp Gly Thr Trp Tyr Ala
210 215 220
Asn Glu Leu Gln Leu Asp Gly Phe Arg Leu Asp Ala Val Lys His Ile
225 230 235 240
Lys Phe Ser Phe Leu Arg Asp Trp Val Asn His Val Arg Glu Lys Thr
245 250 255

EP 0 867 504 A1

Gly Lys Glu Met Phe Thr Val Ala Glu Tyr Trp Gln Asn Asp Leu Gly
 250 265 270
 5 Ala Leu Glu Asn Tyr Leu Asn Lys Thr Asn Phe Asn His Ser Val Phe
 275 280 285
 Asp Val Pro Leu His Tyr Gln Phe His Ala Ala Ser Thr Gln Gly Gly
 290 295 300
 10 Gly Tyr Asp Met Arg Lys Leu Leu Asn Gly Thr Val Val Ser Lys His
 305 310 315 320
 Pro Leu Lys Ser Val Thr Phe Val Asp Asn His Asp Thr Gln Pro Gly
 325 330 335
 15 Gln Ser Leu Glu Ser Thr Val Gln Thr Trp Phe Lys Pro Leu Ala Tyr
 340 345 350
 Ala Phe Ile Leu Thr Arg Glu Ser Gly Tyr Pro Gln Val Phe Tyr Gly
 355 360 365
 20 Asp Met Tyr Gly Thr Lys Gly Asp Ser Gln Arg Glu Ile Pro Ala Leu
 370 375 380
 Lys His Lys Ile Glu Pro Ile Leu Lys Ala Arg Lys Gln Tyr Ala Tyr
 385 390 395 400
 25 Gly Ala Gln His Asp Tyr Phe Asp His His Asp Ile Val Gly Trp Thr
 405 410 415
 Arg Glu Gly Asp Ser Ser Val Ala Asn Ser Gly Leu Ala Ala Leu Ile
 420 425 430
 30 Thr Asp Gly Pro Gly Gly Ala Lys Arg Met Tyr Val Gly Arg Gln Asn
 435 440 445
 Ala Gly Glu Thr Trp His Asp Ile Thr Gly Asn Arg Ser Glu Pro Val
 450 455 460
 35 Val Ile Asn Ser Glu Gly Trp Gly Glu Phe His Val Asn Gly Gly Ser
 465 470 475 480
 Val Ser Ile Tyr Val Gln Arg
 485

(2) INFORMATION FOR SEQ ID NO:38:

40 (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 32 amino acids
 (B) TYPE: amino acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

45 (ii) MOLECULE TYPE: protein

(x1) SEQUENCE DESCRIPTION: SEQ ID NO:38:

50 Met Lys Gln Gln Lys Arg Leu Thr Ala Arg Leu Leu Thr Leu Leu Phe
 1 5 10 15
 Ala Leu Ile Phe Leu Leu Pro His Ser Ala Ala Ala Ala Ala Asn Leu
 20 25 30

55 (2) INFORMATION FOR SEQ ID NO:39:

(i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 33 amino acids
 (B) TYPE: amino acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:39:

Met Arg Ser Lys Thr Leu Trp Ile Ser Leu Leu Phe Ala Leu Thr Leu
 1 5 10 15

Ile Phe Thr Met Ala Phe Ser Asn Met Ser Ala Gln Ala Ala Gly Lys
 20 25 30

Ser

(2) INFORMATION FOR SEQ ID NO:40:

(i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 35 amino acids
 (B) TYPE: amino acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:40:

Met Arg Ser Lys Thr Leu Trp Ile Ser Leu Leu Phe Ala Leu Thr Leu
 1 5 10 15

Ile Phe Thr Met Ala Phe Ser Asn Met Ser Ala Gln Ala Ala Ala Ala
 20 25 30

Ala Ala Asn
 35

(2) INFORMATION FOR SEQ ID NO:41:

(i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 32 amino acids
 (B) TYPE: amino acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:41:

Met Arg Ser Lys Thr Leu Trp Ile Ser Leu Leu Phe Ala Leu Thr Leu
 1 5 10 15

Ile Phe Thr Met Ala Phe Ser Asn Met Ser Ala Gln Ala Ala Asn Leu
 20 25 30

(2) INFORMATION FOR SEQ ID NO:42:

(i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 33 base pairs

(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

5 (ii) MOLECULE TYPE: DNA (genomic)

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:42:

10 CACCTAATTA AAGCTTTCAC ACATTTTCAT TTT 33

(2) INFORMATION FOR SEQ ID NO:43:

(i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 33 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

15

(ii) MOLECULE TYPE: DNA (genomic)

20

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:43:

CACCTAATTA AAGCTTACAC ACATTTTCAT TTT 33

(2) INFORMATION FOR SEQ ID NO:44:

(i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 66 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

25

(ii) MOLECULE TYPE: DNA (genomic)

30

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:44:

35 CCGCGTAATT TCCGGAGAAC ACCTAATTAA AGCCGCAACA CATTTTCATT TTCCCGGGCG 60
CGGCAG 66

(2) INFORMATION FOR SEQ ID NO:45:

(i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 42 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

40

(ii) MOLECULE TYPE: DNA (genomic)

45

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:45:

CCGGAGAACA CCTAATTAA GCCCTAACAC ATTTTCATT TC 42

50

(2) INFORMATION FOR SEQ ID NO:46:

(i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 42 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single

55

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:46:

CCGGAGAACA CCTAATTAAA GCCCACACAC ATTTTCATTT TC

42

(2) INFORMATION FOR SEQ ID NO:47:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 42 base pairs

(B) TYPE: nucleic acid

(C) STRANDEDNESS: single

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:47:

CCGGAGAACA CCTAATTAAA GCCTGCACAC ATTTTCATTT TC

42

(2) INFORMATION FOR SEQ ID NO:48:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 24 base pairs

(B) TYPE: nucleic acid

(C) STRANDEDNESS: single

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:48:

GATGCAGTAT TTCGAACTGG TATA

24

(2) INFORMATION FOR SEQ ID NO:49:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 26 base pairs

(B) TYPE: nucleic acid

(C) STRANDEDNESS: single

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:49:

TGCCCCAATGA TGGCCAACAT TGGAAG

26

(2) INFORMATION FOR SEQ ID NO:50:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 24 base pairs

(B) TYPE: nucleic acid

(C) STRANDEDNESS: single

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:50:

CGAATGGTAT GCTCCCAATG ACGG

24

(2) INFORMATION FOR SEQ ID NO:51:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 24 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:51:

CGAATGGTAT CGCCCAATG ACGG

24

(2) INFORMATION FOR SEQ ID NO:52:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 24 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:52:

CGAATGGTAT AATCCCAATG ACGG

24

(2) INFORMATION FOR SEQ ID NO:53:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 24 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:53:

CGAATGGTAT GATCCCAATG ACGG

24

(2) INFORMATION FOR SEQ ID NO:54:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 24 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:54:

CGAATGGTAT CACCCCAATG ACGG

24

(2) INFORMATION FOR SEQ ID NO:55:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 24 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:55:

CGAATGGTAT AAACCCAATG ACGG

24

(2) INFORMATION FOR SEQ ID NO:56:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 24 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:56:

CGAATGGTAT CCGCCCAATG ACGG

24

(2) INFORMATION FOR SEQ ID NO:57:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 24 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:57:

CGAATGGTAT TCTCCCAATG ACGG

24

(2) INFORMATION FOR SEQ ID NO:58:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 24 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:58:

CGAATGGTAC ACTCCCAATG ACGG

24

(2) INFORMATION FOR SEQ ID NO:59:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 24 base pairs

(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:59:

CGAATGGTAT GTTCCCAATG ACGG

24

(2) INFORMATION FOR SEQ ID NO:60:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 24 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:60:

CGAATGGTAT TGTCCCAATG ACGG

24

(2) INFORMATION FOR SEQ ID NO:61:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 24 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:61:

CGAATGGTAT CAACCCAATG ACGG

24

(2) INFORMATION FOR SEQ ID NO:62:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 24 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:62:

CGAATGGTAT GAACCCAATG ACGG

24

(2) INFORMATION FOR SEQ ID NO:63:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 24 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

5 (xi) SEQUENCE DESCRIPTION: SEQ ID NO:63:

CGAATGGTAT GGTCCCAATG ACGG

24

(2) INFORMATION FOR SEQ ID NO:64:

10 (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 24 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

15 (ii) MOLECULE TYPE: DNA (genomic)

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:64:

20 CGAATGGTAT ATTCCCAATG ACGG

24

(2) INFORMATION FOR SEQ ID NO:65:

25 (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 24 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

30

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:65:

CGAATGGTAT TTTCCCAATG ACGG

24

35 (2) INFORMATION FOR SEQ ID NO:66:

(i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 24 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 40 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

45

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:66:

CGAATGGTAC TGGCCCAATG ACGG

24

(2) INFORMATION FOR SEQ ID NO:67:

50 (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 24 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

55

(ii) MOLECULE TYPE: DNA (genomic)

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:67:

CGAATGGTAT TATCCCAATG ACGG

24

(2) INFORMATION FOR SEQ ID NO:68:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 24 base pairs

(B) TYPE: nucleic acid

(C) STRANDEDNESS: single

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:68:

CCGTCATTGG GACTACGTAC CATT

24

Claims

1. A mutant alpha-amylase that is the expression product of a mutated DNA sequence encoding an alpha-amylase, the mutated DNA sequence being derived from a precursor alpha-amylase which is a *Bacillus* alpha- amylase by substitution or deletion of an aminoacid at the position equivalent to M+15 in *B. licheniformis* alpha-amylase, with the proviso that the substituent amino acid is not Leu, Ile, Asn, Ser, Gln, Asp or Glu.

2. A mutant alpha-amylase of claim 1 further comprising one or more other site specific mutations.

3. A mutant alpha-amylase of any preceding claim wherein the precursor is from a *Bacillus* selected from the group *B. licheniformis*, *B. stearothermophilus* and *B. amyloliquefaciens*.

4. A mutant alpha-amylase of claim 3 wherein the precursor is *Bacillus licheniformis* alpha-amylase.

5. DNA encoding a mutant alpha-amylase of any one of claims 1 to 4.

6. Expression vectors encoding the DNA of claim 5.

7. Host cells transformed with the expression vector of claim 6.

8. A detergent composition comprising a mutant alpha-amylase of any one of claims 1 to 4

9. A detergent composition of claim 8 which is a liquid, gel or granular composition.

10. A detergent composition of claim 8 or claim 9 further comprising one or more additional enzymes.

11. A starch liquefying composition comprising a mutant alpha-amylase of any one of claims 1 to 4.

12. A detergent composition which comprises a mutant alpha-amylase and one or more additional enzymes wherein said mutant alpha-amylase is the expression product of a mutated DNA sequence encoding an alpha-amylase, the mutated DNA sequence being derived from a precursor alpha-amylase which is a *Bacillus* alpha-amylase by substitution or deletion of an amino acid at the position equivalent to M+15 in *B. licheniformis* alpha-amylase.

13. The detergent composition of claim 12 wherein said mutant alpha-amylase is M15L.

14. The detergent composition of claim 12 or claim 13 wherein said mutant alpha-amylase comprises one or more

other site specific mutations.

16. A detergent composition as claimed in any one of claims 13 to 16 wherein said additional enzyme or enzymes is selected from the group consisting of amylases, proteases, lipases and cellulases.

17. A method of liquefying a granular starch slurry from either a wet or dry milling process at a pH of from about 4 to about 6 comprising:

(a) adding an effective amount of an alpha-amylase mutant to the slurry;

(b) optionally adding an effective amount of an antioxidant to the slurry; and

(c) reacting the slurry for an appropriate time and at an appropriate temperature to liquefy the starch;

wherein said alpha-amylase mutant is the expression product of a mutated DNA sequence encoding an alpha-amylase, the mutated DNA sequence being derived from a precursor alpha-amylase which is a *Bacillus* alpha-amylase by substitution or deletion of an amino acid at the position equivalent to M+15 in *B.licheniformis* alpha-amylase.

18. A starch liquefying composition which comprises a mutant alpha-amylase wherein said mutant is the expression product of a mutated DNA sequence encoding an alpha-amylase, the mutated DNA sequence being derived from a precursor alpha-amylase which is a *Bacillus* alpha-amylase, by substitution or deletion of an amino acid at the position equivalent to M+15 in *B. licheniformis* alpha-amylase.

19. The starch liquefying composition of claim 18 wherein said mutant alpha-amylase is M15L.

10 30 50
 AGCTTGAAGAAGTGAAGAAGCAGAGAGGCTATTGAATAAATGAGTAGAAAGCGCCATATC
 70 90 110
 GGCGCTTTTCTTTTGGGAAGAAAATATAGGGGAAAATGGTACTTGTTAAAAATTTCGGAATAT
 130 150 170
 TTATACAACATCATATGTTTCACATTGAAAGGGGAGGAGAATCATGAAACAACAAAAACG
 M K Q Q K R
 190 210 230
 GCTTTACGCCCCGATTGCTGACGCTGTTATTTGCGCTCATCTTCTTGCTGCCTCATTCTGC
 L Y A R L L T L L F A L I F L L P H S A
 250 270 290
 AGCAGCGGCGGCAAATCTTAATGGGACGCTGATGCAGTATTTTGAATGGTACATGCCCAA
 A A A A N L N G T L M O Y F E W Y M P N
 310 330 350
 TGACGGCCAACATTGGAAGCGTTTGCAAAACGACTCGGCATATTTGGCTGAACACGGTAT
 D G O H W K R L O N D S A Y L A E H G I
 370 390 410
 TACTGCCGTCTGGATTCCCCCGGCATATAAGGGAACGAGCCAAGCGGATGTGGGCTACGG
 T A V W I P P A Y K G T S O A D V G Y G
 430 450 470
 TGCTTACGACCTTTATGATTTAGGGGAGTTTCATCAAAAAGGGACGGTTCGGACAAAGTA
 A Y D L Y D L G E F H Q K G T V R T K Y
 490 510 530
 CGGCACAAAAGGAGAGCTGCAATCTGCGATCAAAAGTCTTCATTCCCGCGACATTAACGT
 G T K G E L O S A I K S L H S R D I N V
 550 570 590
 TTACGGGGATGTGGTCATCAACCACAAAGGCGGCGCTGATGCGACCGAAGATGTAACCGC
 Y G D V V I N H K G G A D A T E D V T A
 610 630 650
 GGTTGAAGTCGATCCCGCTGACCGCAACCGCGTAATTTTCAAGGAGAACACCTAATTAAAGC
 V E V D P A D R N R V I S G E H L I K A
 670 690 710
 CTGGACACATTTTCATTTTCCGGGGCGCGGCAGCACATACAGCGATTTTAAATGGCATTG
 W T H F H F P G R G S T Y S D F K W H W
 730 750 770
 GTACCATTTTGACGGAACCGATTGGGACGAGTCCCGAAAGCTGAACCGCATCTATAAGTT
 Y H F D G T D W D E S R K L N R I Y K F
 790 810 830
 TCAAGGAAAGGCTTGGGATTGGGAAGTTTCCAATGAAAACGGCAACTATGATTATTTGAT
 Q G K A W D W E V S N E N G N Y D Y L M

FIG. 1A

850 870 890
 GTATGCCGACATCGATTATGACCATCCTGATGTCGCAGCAGAAATTAAGAGATGGGGCAC
 Y A D I D Y D H P D V A A E I K R W G T
 910 930 950
 TTGGTATGCCAATGAACTGCAATTGGACGGTTTCCGTCTTGATGCTGTCAAACACATTAA
 W Y A N E L Q L D G F R L D A V K H I K
 970 990 1010
 ATTTTCTTTTTTGC GGGATTGGGTTAATCATGTCAGGGAAAAACGGGGAAGGAAATGTT
 F S F L R D W V N H V R E K T G K E M F
 1030 1050 1070
 TACGGTAGCTGAATATTGGCAGAATGACTTGGGCGCTCTGGAAAACTATTTGAACAAAAC
 T V A E Y W O N D L G A L E N Y L N K T
 1090 1110 1130
 AAATTTTAATCATTCACTGTTTGACGTGCCGCTTCATTATCAGTTCCATGCTGCATCGAC
 N F N H S V F D V P L H Y O F H A A S T
 1150 1170 1190
 ACAGGGAGGCGGCTATGATATGAGGAAATTGCTGAACGGTACGGTCGTTTCCAAGCATCC
 O G G G Y D M R K L L N G T V V S K H P
 1210 1230 1250
 GTTGAAATCGGTTACATTTGTCGATAACCATGATACACAGCCGGGGCAATCGCTTGAGTC
 L K S V T F V D N H D T Q P G O S L E S
 1270 1290 1310
 GACTGTCCAAACATGGTTTAAGCCGCTTGCTTACGCTTTTATTCTCACAAGGGAATCTGG
 T V Q T W F K P L A Y A F I L T R E S G
 1330 1350 1370
 ATACCCTCAGGTTTTCTACGGGGATATGTACGGGACGAAAGGAGACTCCCAGCGCGAAAT
 Y P O V F Y G D M Y G T K G D S O R E I
 1390 1410 1430
 TCCTGCCTTGAAACACAAAATTGAACCGATCTTAAACGCAGAAAACAGTATGCGTACGG
 P A L K H K I E P I L K A R K O Y A Y G
 1450 1470 1490
 AGCACAGCATGATTATTTGACCAACCATGACATTGTCGGCTGGACAAGGGAAGGCGACAG
 A O H D Y F D H H D I V G W T R E G D S
 1510 1530 1550
 CTCGGTTGCAAATTCAGGTTTGGCGGCATTAATAACAGACGGACCCGGTGGGGCAAAGCG
 S V A N S G L A A L I T D G P G G A K R
 1570 1590 1610
 AATGTATGTCGGCCGGCAAACGCGGTGAGACATGGCATGACATTACCGGAAACCGTTC
 M Y V G R C N A G E T W H D I T G N R S
 1630 1650 1670
 GGAGCCGGTTGTCATCAATTTCGGAAGGCTGGGGAGAGTTTCACGTAAACGGCGGGTCGGT
 E P V V I N S E G W G E F H V N G G S V

FIG. 1B

1690 1710 1730
 TTCAATTTATGTTCAAAGATAGAAGAGCAGAGAGGACGGATTTCTGAAGGAAATCCGTT
 S I Y V Q R .
 1750 1770 1790
 TTTTATTTTGCCCGTCTTATAAATTTCTTTGATTACATTTTATAATTAATTTTAACAAA
 1810 1830 1850
 GTGTCATCAGCCCTCAGGAAGGACTTGCTGACAGTTTGAATCGCATAGGTAAGGCGGGGA
 1870 1890 1910
 TGAAATGGCAACGTTATCTGATGTAGCAAAGAAAGCAAATGTGTGCGAAAATGACGGTATC
 1930 1950
 GCGGGTGATCAATCATCCTGAGACTGTGACGGATGAATTGAAAAAGCT

FIG._1C




FIG._1

10 30 50
ANLNGTLMQYFEWYMPNDGOHWKRLQND SAYLAEHGITAVWIPPAYKGT SOADVGYGAYD

70 90 110
LYDLGEFHQKGTVRTKYGTKGELOSAIKSLHSRDINVYGDVVINHKGGADATEDVTAVEV

130 150 170
DPADRN RVISGEHLIKAWTHFHFPGRGSTYSDFKWHWYHFDGTDWDESRKLNRIYKFOGK

190 210 230
AWDWEVSNENGNYDYLMYADIDYDHPDVAAEIKRWGTWYANELQLDGFRLDAVKHIKFSF

250 270 290
LRDWVNHVREKTGKEMFTVAEYWONDLGALENYLNKTNFNHSVFDVPLHYOFHAASTOGG

310 330 350
GYDMRKLLNGTVVSKHPLKSVTFVDNHDTOPGOSLESTVOTWFKPLAYAFILTRESGYPO

370 390 410
VFYGD MYGTKGDSOREIPALKHKIEPILKARKOYAYGAQH DYFDHHDIVGWTREGDSSVA

430 450 470
NSGLAALITDGP GGAKRM YVGRONAGETWHDITGNRSEPVVINSEGWGEFHVNGG SVSIY

VQR


FIG. 2

Am-Lich = <i>B. Licheniformis</i>	Am-Amylo = <i>B. amyloliquefaciens</i>	Am-Stearo = <i>B. stearothermophilus</i>
1 Am-Lich Am-Amylo Am-Stearo	1MKQQ KRLYARLLTL LFALIFLLPH MRGRGNMIQK RKRTVSFRLV LMCTLLFVSLVLTF HRIIRKGMWF LLAFLLTASL	1SAAA AANLNGTLMQ YFEWYMPNDGPITK TSAVNGTLMQ YFEWYTPNDG FCPTGRHAKA AAPFNGTMMQ YFEWYLPDDG
61 Am-Lich Am-Amylo Am-Stearo	61 QIIWKRLQND S AYLAEHGITA VWIPPAYKGT QIIWKRLQND A LILSDIGITA VWIPPAYKGL ILWTKVANE A NNLSLIGITA LSLPPAYKGL	79 120 KGTVRIKYGT KGTVRIKYGT KGTVRIKYGT
121 Am-Lich Am-Amylo Am-Stearo	121 KGELQSAIKS LHSRDINVYG DVVINIKGGA DATEDVTAVE KSELQDAIGS LHSRNVQVYG DVVLNHHKAGA DATEDVTAVE KAQYLOAIQA AHAAGMOVYA DVVFDHKGGA DGTEWVDAVE	139 180 SGEHLIKAWT SEEYQIKAWT SGTYQIQAWT
181 Am-Lich Am-Amylo Am-Stearo	181 IIFHFPGRGST YSDFKWHWYH FDGTDWDESR KLNRIYKF.... DFRFPGRGNT YSDFKWHWYH FDGADWDESR KISRIFKFRG KDFPGRGNT YSDFKWRWYH FDGVDWDESR KLSRIYKFRG	197 240 NENGNYDYLM SENGNYDYLM TENGNYYDYM
241 Am-Lich Am-Amylo Am-Stearo	241 YADIDYDHPD VAAEIKRWGT WYANELQLDG FRLDAVKHIK YADVVDYDHPD VVAETKKWGI WYANELSDLG FRIDAAKHIK YADLMDHPE VVTELKNWVK WYVNTTNIDG <u>FRLDGLKHIK</u>	257 300 VREKTGKEMF VRQATGKEMF VRSQTGKPLF
301 Am-Lich Am-Amylo Am-Stearo	301 TVAEYWQNDL GALENYLNKT NFNHVSFVDP LHYQFHAAS TVAEYWQNN A GKLENYLNKT SFNQSVFVDP LHFNLQAASS TVGEYWSYDI <u>NKLHNYITKT</u> NGTMSLFDAP LHNKFYTASK	317 360 OGGGYDMRKL OGGGYDMRRL SGGAFDMRTL

FIG. 3A

Am-Lich	361	LKSVTFVDNH	DTQPGQSLES	TVQTWFKPLA	YAFILTRESG	YQVIFYGDMY	GTKGDSQREI	377
Am-Amylo		EKAVTFVENH	DTQPGQSLES	TVQTWFKPLA	YAFILTRESG	YQVIFYGDMY	GTKGTSPKEI	420
Am-Stearo		TLAVTFVDNH	DTNPAKR.CS	HGRPWFKPLA	YAFILTROEG	YPCVIFYGDYY	GI.....POYNI	
Am-Lich	421	PALKKKIEPI	LKARKQYAYG	AQHDIYTDIHD	IVGWTREGDS	SVANSGLAAL	ITDGPGGAKR	437
Am-Amylo		PSLKDNIEPI	LKARKEYAYG	POHDYIDIHPD	VIGWTREGDS	SAKSGLAAL	ITDGPGGSKR	480
Am-Stearo		PSLKSKIDPL	LIARRDYAYG	TQHDYLDHSD	IIGWTREGVI	EKPGSGLAAL	ITDGAGRSKW	
Am-Lich	481	MYVGRQNAGE	TWHDITGNRS	EPVVINSEGW	GEFHVNGGSV	SIYVQR.....		483
Am-Amylo		MYAGLKNAGE	TWYDITGNRS	DTVKIGSDGW	GEFHVNDGSV	SIYVQK.....		540
Am-Stearo		MYVKGQHAGK	VFYDLTGNRS	DTVTINSDGW	GEFKVNGGSV	SVWVPRKTTV	STIARPITTR	
Am-Lich	541	
Am-Amylo		
Am-Stearo		PWTGEEFVRWH	EPRLVAWP.	
				559				

FIG._3B



10 30 50
 ANLNGTLMQYFEWYMPNDGOHWKRLONDSAYLAEHGITAVWIPPAYKGT SQADVGYGAYD
 70 90 110
 LYDLGEFHQKGTVRTKYGTKGELOSAIKSLHSRDINVYGDVVINHKGGADATEDVTAVEV
 130 150 170
 DPADRN RVISGEHLIKAWTHFHFPGRGSTYSDFKWHWYHFDGTDWDESRKLNRIYKFOGK
 190 210 230
 AWDWEVSNENGNYDYLT₁YADIDYDHPDVAAEIKRWGTWYANELQLDGFRLDAVKHIKFSF
 250 270 290
 LRDWVNHVREKTGKEMFTVAEYWONDLGALENYLNKTNFNHVSFVDVPLHYQFHAASTQGG
 310 330 350
 GYDMRKLLNGTVVSKHPLKSVTFVDNHDTQPGOSLESTVOTWFKPLAYAFILTRESGYPO
 370 390 410
 VFYGD MYGTKGDSOREIPALKHKIEPILKARKOYAYGAOH DYFDHHDIVGWTREGDSSVA
 430 450 470
 NSGLAALITDGP GGAKRMYVGRONAGETWHDITGNRSEPVVINSEGWGEFHVNGGSVSIY
 VQR

FIG. 4a

AAAA

14 34 54
ANLNGTLMQYFEWYMPNDGOHWKRLONDSAYLAEHGITAVWIPPAYKGT SQADVG YGAYD

74 94 114
LYDLGEFHQKGTVRTKYGTKGELOSAIKSLHSRDINVYGDVVINHKGGADATEDVTAVEV

134 154 174
DPADRN RVISGEHLIKAWTHFHFPGRGSTYSDFKWHWYHFDGTDWDESRKLNRIYKFQ GK

194 214 234
AWDWEVSNENGNYDYLMYADIDYDHPDVAAEIKRWGTWYANELQLDGFRLDAVKHIKFSF

254 274 294
LRDWVNVHVREKTGKEMFTVAEYWONDLGALENYLNKTNFNHVSF DVPLHYOFHAASTQGG

314 334 354
GYDMRKLLNGTVVSKHPLKSVTFVDNHDTPGQSLESTVQTWFKPLAYAFILTRESGYPO

374 394 414
VFYGD MYGTKGDSOREIPALKHKIEPILKARKOYAYGAOHDYFDHHDIVGWTREGDSSVA

434 454 474
NSGLAALITDGP GGAKRMYVGRONAGETWHDITGNRSEPVVINSEGWGEFHVNGGSVSIY

VQR

FIG. 4b




FIG. 5

SIGNAL SEQUENCE - MATURE PROTEIN JUNCTIONS IN:

B.licheniformis alpha-amylase.

MKQOKRLTARLLTLLFALIFLLPHSA^(PstI)AAA[ANL.....
N-terminus

B.subtilis alkaline protease aprE.

MRSKTLWISLLFALTLLFTMAFSNMSAOA^(PstI)AGKS.....
N-terminus

B.licheniformis alpha-amylase in pA4BL.

MRSKTLWISLLFALTLLFTMAFSNMSAOA^(PstI)AAAAAN.
N-terminus

B.licheniformis alpha-amylase in pBLapr.

MRSKTLWISLLFALTLLFTMAFSNMSAOA[ANL.....
N-terminus

(PstI) indicates the site of the restriction site in the gene.


↓

[

N-terminus indicates cleavage site between signal peptide and secreted protein.

FIG._6

FIG. 7



11/22

FIG. 8




FIG. 9




FIG. 10




FIG. 12




FIG. 11




FIG. 13




FIG. 15





FIG. 14





European Patent
Office

EUROPEAN SEARCH REPORT

Application Number
EP 98 10 9967.4

DOCUMENTS CONSIDERED TO BE RELEVANT			
Category	Citation of document with indication, where appropriate, of relevant passages	Relevant to claim	CLASSIFICATION OF THE APPLICATION (Int. Cl.4)
Y	THE JOURNAL OF BIOLOGICAL CHEMISTRY, Volume 260, No 11, 1985, David A. Estell et al, "Engineering an Enzyme by Site-directed Mutagenesis to Be Resistant to Chemical Oxidation" * page 6518 - page 6521 see page 6518, line 1 - line 10 *	1-19	C12N 9/28 C12N 15/56 C11D 3/386
Y	EP 0410498 A2 (GIST-BROCADES N.V.), 30 January 1991 (30.01.91) * page 5, line 19 - line 20; page 6, line 35 - line 40 *	1-19	
A	WO 9116423 A1 (NOVO NORDISK A/S), 31 October 1991 (31.10.91) * claims 1-2 *	1-19	
A	BIOTECHNOLOGY, Volume 10, December 1992, Philippe Joyet et al, "Hyperthermostable variants of a highly thermostable alpha-amylase" * page 1579 - page 1583 figure 4 *	1-19	
The present search report has been drawn up for all claims			TECHNICAL FIELDS SEARCHED (Int. Cl.6)
			C12N
Place of search		Date of completion of the search	Examiner
STOCKHOLM		16 July 1998	PATRICK ANDERSSON
<p>CATEGORY OF CITED DOCUMENTS</p> <p>X : particularly relevant if taken alone Y : particularly relevant if combined with another document of the same category A : technological background O : non-written disclosure P : intermediate document</p> <p>T : theory or principle underlying the invention E : earlier patent document, but published on or after the filing date D : document cited in the application I : document cited for other reasons A : member of the same patent family, corresponding document</p>			

BEST AVAILABLE COPY

EP 0 867 504 A1 (1998)